

# Interventions to optimize embryo transfer in women undergoing assisted conception: a comprehensive systematic review and meta-analyses

Bede Tyler<sup>1</sup>, Hugo Walford<sup>1</sup>, Jennifer Tamblyn<sup>2</sup>, Stephen D. Keay<sup>3</sup>, Dimitrios Mavrelas<sup>1,4</sup>, Ephia Yasmin<sup>1,4</sup>, and Bassel H. Al Wattar <sup>1,4,\*</sup>

<sup>1</sup>UCL Institute for Women's Health, University College London, London, UK <sup>2</sup>Institute of Metabolism and Systems Research (IMSR), University of Birmingham, Birmingham, UK <sup>3</sup>Centre for Reproductive Medicine, University Hospital of Coventry & Warwickshire, Coventry, UK <sup>4</sup>Reproductive Medicine Unit, Elizabeth Garrett Anderson Wing, University College London Hospitals, London, UK

\*Correspondence address. Reproductive Medicine Unit, Elizabeth Garrett Anderson Wing, University College London Hospitals, London, UK, WC1E 6DB. E-mail: [dr.basselwa@gmail.com](mailto:dr.basselwa@gmail.com)  <https://orcid.org/0000-0001-8287-9271>

Submitted on October 15, 2021; resubmitted on February 2, 2022; editorial decision on February 10, 2022

---

## TABLE OF CONTENTS

---

- Introduction
  - Methods
    - Search strategy
    - Review selection and inclusion process
    - Data extraction
    - Quality assessment and confidence of evidence
    - Statistical analysis
  - Results
    - Quality assessment
    - Pharmacological interventions
    - Embryo transfer techniques
    - Relaxation interventions
  - Discussion
    - Summary of main findings
    - Strengths and limitations
    - Implications for clinical practice
    - Future research need
  - Conclusion
- 

**BACKGROUND:** Several interventions and techniques are suggested to improve the outcome of embryo transfer (ET) in assisted conception. However, there remains no consensus on the optimal practice, with high variations among fertility specialists.

**OBJECTIVE AND RATIONALE:** We conducted a comprehensive systematic review and meta-analyses of randomized controlled trials (RCTs) aiming to identify effective interventions that could be introduced around the time of ET to improve reproductive outcomes.

**SEARCH METHODS:** We searched the electronic databases (MEDLINE, EMBASE and Cochrane CENTRAL) from inception until March 2021 using a multi-stage search strategy of MeSH terms and keywords, and included all RCTs that evaluated an intervention in the 24-h period before/after ET in women undergoing IVF/ICSI. Our primary outcome was clinical pregnancy rate post-ET confirmed as viable pregnancy on ultrasound scan. We assessed the risk of bias in included trials and extracted data in duplicate. We pooled data using a random-effect meta-analysis and reported using risk ratio (RR) with 95% CI. We explored publication bias and effect modifiers using subgroup analyses.

**OUTCOMES:** Our search yielded 3685 citations of which we included 188 RCTs (38 interventions, 59 530 participants) with a median sample size of 200 (range 26–1761). The quality of included RCTs was moderate with most showing a low risk of bias for randomization (118/188, 62.8%) and attrition (105/188, 55.8%) but there was a significant risk of publication bias (Egger's test  $P=0.001$ ). Performing ET with ultrasound guidance versus clinical touch ( $n=24$ , RR 1.265, 95% CI 1.151–1.391,  $I^2=38.53\%$ ), hyaluronic acid versus routine care ( $n=9$ , RR 1.457, 95% CI 1.197–1.261,  $I^2=46.48\%$ ) and the use of a soft versus hard catheter ( $n=27$ , RR 1.122, 95% CI 1.028–1.224,  $I^2=57.66\%$ ) led to higher clinical pregnancy rates. Other pharmacological add-ons also showed a beneficial effect including granulocyte colony-stimulating factor (G-CSF:  $n=4$ , RR 1.774, 95% CI 1.252–2.512,  $I^2=0$ ), Atosiban ( $n=7$ , RR 1.493, 95% CI 1.184–1.882,  $I^2=68.27\%$ ) and hCG ( $n=17$ , RR 1.232, 95% CI 1.099–1.382,  $I^2=57.76\%$ ). Bed rest following ET was associated with a reduction in clinical pregnancy ( $n=6$ , RR 0.857, 95% CI 0.741–0.991,  $I^2=0.01\%$ ). Other commonly used interventions, such as non-steroidal anti-inflammatory drugs, prophylactic antibiotics, acupuncture and cervical mucus removal, did not show a significant benefit on reproductive outcomes. Our effect estimates for other important outcomes, including miscarriage and live birth, were limited by the varied reporting across included RCTs.

**WIDER IMPLICATIONS:** Using ultrasound guidance, soft catheters and hyaluronic acid at the time of ET appears to increase clinical pregnancy rates. The use of Atosiban, G-CSF and hCG showed a trend towards increased clinical pregnancy rate, but larger trials are required before adopting these interventions in clinical practice. Bed rest post-ET was associated with a reduction in clinical pregnancy and should not be recommended.

**Key words:** embryo transfer / assisted conception / IVF / ICSI / systematic review / meta-analysis

## Introduction

Rapid developments in ART offered hope to many subfertile couples, with more than 8 million ART babies conceived worldwide (Steptoe and Edwards, 1978; Zegers-Hochschild *et al.*, 2009; Cardona Barberán *et al.*, 2020). ART treatments are complex, comprising several interrelated steps often with high stress and psychological bearings on couples under treatment (Malina and Pooley, 2017). Numerous interventions or 'add-on' therapies have been introduced over the last few decades to optimize the outcome of ART treatments and assisted conception rates (Farquhar, 2019). However, the effectiveness and safety of several add-ons remain unclear raising concerns about patients' safety and the vulnerability to profiteering (Galiano *et al.*, 2021).

Embryo transfer (ET) is a crucial component to the success of ART treatments (Mains and van Voorhis, 2010). While seemingly a simple procedure, ET is operator dependent with suboptimal practice often linked to cycle failure and reduced pregnancy rates (Yao *et al.*, 2009a). Some add-ons are proposed at the time of ET to increase the chances of conception, such as the use of ultrasound guidance (Cozzolino *et al.*, 2018), pharmacological interventions aimed to minimize uterine contractility at the time of the transfer (Ng *et al.*, 2019; Schwarze *et al.*, 2020) and also pre-transfer acupuncture and relaxation techniques (Smith *et al.*, 2019). Currently, the practice of ET varies among fertility specialists with no consensus on the optimal ET technique (Nancarrow *et al.*, 2021).

The American Society for Reproductive Medicine (ASRM) guideline recommends several ET techniques and add-ons including the use of ultrasound guidance, soft catheters and avoidance of bed rest (Penzias *et al.*, 2017a). Since its publication in 2016, several new add-ons have been proposed, with more than 20 relevant randomized controlled trials (RCTs) reported. Contemporary evidence synthesis is, therefore,

needed to inform and update the current knowledge gap on ET practice.

To address this research need, we conducted a comprehensive systematic review and meta-analyses of randomized trials evaluating any intervention introduced at the time of ET to improve reproductive outcomes in couples undergoing ART.

## Methods

We conducted a systematic review using a prospectively registered protocol (PROSPERO: CRD42020216199) and reported in line with established guidelines (Page *et al.*, 2021).

### Search strategy

We searched the electronic databases (MEDLINE, EMBASE and Cochrane CENTRAL) from inception until March 2021 for all RCTs that evaluated an intervention introduced at the time of ET in women undergoing ART treatment. We used a multi-stage search strategy using MeSH terms and keywords, and combined them using the Boolean operators AND/OR to identify relevant citations (Supplementary Data). We did not apply any search filters or language restrictions. We screened the bibliographies of relevant articles and performed complementary searches in Google Scholar and Scopus to identify any missed citations and grey literature.

### Review selection and inclusion process

Two authors (B.T. and H.W.) independently screened the titles and abstracts to identify relevant citations. Full-text articles were then screened against our inclusion criteria. Any discrepancies were resolved through consultation with the senior author (B.H.A.). We

included all RCTs that evaluated any clinical intervention introduced at the time of ET (within 24 h of the procedure) following any ART treatment (IVF and/or ICSI) with the aim to improve implantation rate irrespective of the cause of subfertility. Interventions were defined as 'add-ons' if they were introduced as supplementary to a standardized ART protocol in both groups of comparisons in the 24-h period before/after ET. Comparison groups in included trials received the same ART protocol versus control with no additional treatment (standard care), placebo or sham treatment.

Studies reporting intra-uterine sperm injection or ovulation induction treatments were excluded. Articles not in English were translated and included if deemed relevant. We also excluded quasi, non-randomized, observational and animal studies. Review articles and RCTs that did not report on any reproductive outcomes were also excluded.

## Data extraction

Data extraction was performed in duplicate (B.T. and H.W.) using a piloted electronic data collection tool with the following characteristics collected: study publication year and journal, inclusion–exclusion criteria, type of intervention and comparison evaluated, characteristics of the included study population and the evaluated ART treatments and all reproductive outcomes.

Our primary outcome was clinical pregnancy rate post-ET confirmed as viable pregnancy on ultrasound scan. We also reported live birth, ongoing pregnancy (a viable pregnancy beyond 12 weeks of gestation) and on biochemical pregnancy (confirmed with a positive  $\beta$ hCG test).

## Quality assessment and confidence of evidence

The quality of published literature was assessed by two independent reviewers (B.T. and H.W.) using the Cochrane Risk of Bias assessment tool (ROB2; Sterne et al., 2019). Studies were assessed in five domains: randomization process (randomization bias), deviations from intended interventions (allocation bias), performance bias, missing outcome data (attrition bias), measurement of the outcome (detection bias) and selection of the reported result (reporting bias). Each domain was assessed to have a high, medium or low risk of bias. Due to the nature of the intervention, we did not penalize unblinded trials but reported on assessors blinding. In case of uncertainty, consensus was established with input from a third reviewer (J.T.).

We employed the Grading of Recommendations Assessment, Development and Evaluation (GRADE) approach (GRADE Working Group, 2004) to evaluate the confidence in available evidence for each intervention and aid translation into clinical practice. Two independent reviewers (H.W. and J.T.) assessed the certainty of the evidence for the following domains: risk of bias in primary studies for each intervention, inconsistency across included trials, indirectness or imprecision in the pooled effect estimates. A final overall assessment was made for each intervention as high, moderate, low or very low in consultation with a third reviewer (B.H.A.).

## Statistical analysis

We reported on dichotomous outcomes using summary risk ratio (RR) with 95% CI and continuous outcomes using weighted mean

difference. We pooled data using a random-effect meta-analysis and applied a restricted maximum likelihood model (Lin et al., 2013). Study heterogeneity among included trials was assessed using the  $I^2$  statistics. We also assessed the publication bias and small study effect using a funnel plot for each pairwise comparison and performed Egger's test to assess its statistical significance (Sterne and Harbord, 2004). We planned a sensitivity meta-regression and subgroup analyses to investigate potential effect modifiers where relevant (Harbord and Higgins, 2009). All statistical analyses were conducted in Stata V13 (StataCorp, TX, USA) and Open Meta-analyst software (Brown University; Providence, RI, USA).

## Results

Our electronic search identified 3685 titles and abstracts of which we screened 228 articles in full against our inclusion criteria. We identified a further 65 additional records from screening bibliographies and supplementary searches. In total, we included 188 RCTs reporting on 59 530 participants (Fig. 1). The median sample size across the studies was 200 (range 26–1761). About half of the included studies were conducted in European countries (62/188, 33%) followed by the USA (23/188, 12.2%) and Iran (17/188, 9%). More than a third of included studies performed an ET at Day 2–3 (cleavage stage; 76/188, 40.4%), 8.5% at Day 5–6 (blastocyst; 16/188, 8.5%) and a third did not specify (69/188, 36.7%). Most studies performed a fresh ET (96/188, 51.1%), about a 10th performed either a frozen or fresh ET (21/188, 11.2%; Supplementary Tables SI and SII).

## Quality assessment


Overall, the quality of included RCTs was moderate with most studies assessed as low risk of bias for randomization (118/188, 62.8%) and attrition (105/188, 56.4%). Almost half of the studies were at low risk of bias for outcome reporting (85/188, 45.2%) and one-third were low risk for allocation (58/188, 30.9%) and detection bias (62/188 33.0%). Performance bias was judged to be high in one-third of the included studies (54/188, 28.7%; Fig. 2).

Our funnel plots showed significant variations in effect size and standard error across included RCTs with evidence of small study effect (Fig. 3). We explored this further by removing interventions that were evaluated by a single RCT, which reduced the distribution of the funnel plot (Supplementary Fig. S1). Egger's test showed significant publication bias with a  $P$ -value of 0.001.


## Pharmacological interventions

### Antibiotics

We identified two RCTs that evaluated the effects of using antibiotics at the time of ET (Peikrishvili et al., 2004; Brook et al., 2006;  $n = 625$  participants). The pooled effect estimate showed no significant benefit for the use of antibiotics compared to standard care on the rate of clinical pregnancy (RR 1.008, 95% CI 0.812–1.251,  $I^2 = 0$ ; Supplementary Fig. S2a). Ongoing pregnancy and miscarriage were reported only in one study (Peikrishvili et al., 2004) showing no significant benefit (RR 0.840, 95% CI 0.570–1.239 and RR 1.255, 95% CI 0.499–3.157, respectively).




**Figure 1.** Selection and inclusion process for randomized controlled trials evaluating interventions at the time of embryo transfer in women undergoing assisted reproduction.



**Figure 2.** Risk of bias in included randomized controlled trials evaluating interventions at the time of embryo transfer in women undergoing assisted reproduction.

Atosiban

Seven RCTs assessed the effectiveness of Atosiban to promote uterine relaxation at the time of ET (Ahn *et al.*, 2009; Moraloglu *et al.*, 2010; Song *et al.*, 2013; Ng *et al.*, 2014; He *et al.*, 2016; Hebisha *et al.*, 2016; Yuan *et al.*, 2019) using varied doses and preparations (n= 1646 participants). The pooled effect estimate showed a significant increase in



clinical pregnancy with the use of Atosiban compared to placebo (RR 1.493, 95% CI 1.184–1.882,  $I^2 = 68.27\%$ ; [Supplementary Fig. S3a](#)).

Ongoing pregnancy was shown to increase with the use of Atosiban in three RCTs (RR 1.137, 95% CI 1.007, 1.283,  $P = 0.435$ ,  $I^2 = 0\%$ ; [Ng et al., 2014](#); [Visnova et al., 2018](#); [Bosch et al., 2019](#)) while there was no significant effect on live birth ( $n = 1$ , RR 1.05, 95% CI 0.879–1.240; [Ng et al., 2014](#)), miscarriage ( $n = 5$ , RR 1.080, 95% CI 0.748–1.561,  $I^2 = 0\%$ ; 24, 26–29) or biochemical pregnancy ( $n = 2$ , RR 1.741, 95% CI 0.664–4.567,  $I^2 = 92.92\%$ ; 27, 29; [Supplementary Fig. S3b–d](#)).

A subgroup analysis showed a persistent improvement in clinical pregnancy associated with the use of a medium dose Atosiban (35–50 mg;  $n = 5$ , RR 1.591, 95% CI 1.110–1.714,  $I^2 = 76.68\%$ ; [Ahn et al., 2009](#); [Moraloglu et al., 2010](#); [Song et al., 2013](#); [Ng et al., 2014](#); [Yuan et al., 2019](#)) and as well as a low dose (5–10 mg;  $n = 2$ , RR 1.380, 95% CI 1.110–1.714,  $I^2 = 68.27\%$ ; [He et al., 2016](#); [Hebisha et al., 2016](#); [Supplementary Fig. S3e](#)).

#### Nifedipine

Only one study with low risk of bias evaluated the use of Nifedipine for uterine relaxation at the time of ET ([Ng et al., 2019](#)). There was no improvement in clinical pregnancy (RR 1.115, 95% CI 0.548–2.267), live birth (RR 1.115, 95% CI 0.548–2.267) or miscarriage (RR 1.022, 95% CI 0.105–6.951) with using Nifedipine compared to placebo.

#### Non-steroidal anti-inflammatory drugs

We identified seven studies that evaluated the use of different non-steroidal anti-inflammatory drugs (NSAIDs) at the time of ET, including Piroxicam ( $n = 4$ ; [Moon et al., 2004](#); [Dehghani Firouzabadi et al., 2006](#); [Prato and Borini, 2009](#); [Zarei et al., 2021](#)), Indomethacin ( $n = 1$ ; [Bernabeu et al., 2006](#)), Aspirin ( $n = 1$ ; [Duvan et al., 2006](#)) and Ibuprofen ( $n = 1$ ; [Fekih et al., 2013](#)). A pooled effect estimates for all types of NSAIDs ( $n = 1207$  participants) showed no significant effect on clinical pregnancy (RR 1.294, 95% CI 0.973–1.721,  $I^2 = 63.92\%$ ;

[Supplementary Fig. S4a](#)), miscarriage ( $n = 4$ , RR 0.787, 95% CI 0.391–1.582,  $I^2 = 0\%$ ; [Dehghani Firouzabadi et al., 2006](#); [Prato and Borini, 2009](#); [Fekih et al., 2013](#); [Zarei et al., 2021](#)) and biochemical pregnancy ( $n = 5$ , RR 1.032, 95% CI 0.840–1.268,  $I^2 = 26.69\%$ ; [Bernabeu et al., 2006](#); [Dehghani Firouzabadi et al., 2006](#); [Duvan et al., 2006](#); [Prato and Borini, 2009](#); [Zarei et al., 2021](#); [Supplementary Fig. S4b](#) and c). Subgroup meta-analyses per type of NSAIDs showed similar results with no overall benefit ([Supplementary Fig. S4d](#)).

#### Steroids

Only one study evaluated the effect of Prednisone on reproductive outcomes at the time of ET ([Duvan et al., 2006](#)) which had no significant effect on clinical pregnancy (RR 1.064, 95% CI 0.484–2.337).

#### hCG

Supplementing hCG at the time of ET was the most commonly evaluated pharmacological intervention in 17 RCTs, given as an intrauterine infusion with or without embryo culture media ([Mansour et al., 2011](#); [Cambiaghi et al., 2013](#); [Leao et al., 2013](#); [HoNg et al., 2014](#); [Santibañez et al., 2014](#); [Singh and Singh, 2014](#); [Zarei et al., 2014](#); [Aaleysin et al., 2015](#); [Wirleitner et al., 2015a,b](#); [Dehghani Firouzabadi et al., 2016](#); [Eskandar et al., 2016](#); [Hosseini et al., 2016](#); [Mostajeran et al., 2017](#); [Hafezi et al., 2018](#); [Laokirkkiat et al., 2019](#); [WaNg et al., 2019](#); [Supplementary Fig. S5a](#)). The pooled data showed a significant increase in clinical pregnancy with hCG use ( $n = 17$ , RR 1.232, 95% CI 1.099–1.382,  $I^2 = 57.76\%$ ).

We pooled data from nine RCTs that evaluated the provision of intrauterine hCG compared to standard care, which showed a significant improvement in clinical pregnancy with hCG use ( $n = 9$ , RR 1.269, 95% CI 1.092–1.474,  $I^2 = 43.92\%$ ; [Mansour et al., 2011](#); [Cambiaghi et al., 2013](#); [Leao et al., 2013](#); [Santibañez et al., 2014](#); [Dehghani Firouzabadi et al., 2016](#); [Eskandar et al., 2016](#); [Hosseini et al., 2016](#); [Mostajeran et al., 2017](#); [WaNg et al., 2019](#)). Eight of the included studies compared hCG injections to a controlled intrauterine injection of embryo culture medium, which did not show a significant improvement in clinical pregnancy rates ( $n = 8$ , 1.193, 95% CI 0.996–1.428,  $I^2 = 69.7\%$ ; [HoNg et al., 2014](#); [Singh and Singh, 2014](#); [Zarei et al., 2014](#); [Aaleysin et al., 2015](#); [Wirleitner et al., 2015a,b](#); [Hafezi et al., 2018](#); [Laokirkkiat et al., 2019](#); [Supplementary Fig. S5g](#)).

Overall, hCG supplementation did not significantly increase live birth ( $n = 7$ , RR 1.132, 95% CI 0.924–1.387,  $I^2 = 73.37\%$ ; [Supplementary Fig. S5e](#); [Mansour et al., 2011](#); [Singh and Singh, 2014](#); [Aaleysin et al., 2015](#); [Wirleitner et al., 2015a,b](#); [Hafezi et al., 2018](#); [Laokirkkiat et al., 2019](#)), ongoing pregnancy ( $n = 2$ , RR 1.339, 95% CI 0.947–1.893,  $I^2 = 78.24\%$ ; [Supplementary Fig. S5c](#); [HoNg et al., 2014](#); [Aaleysin et al., 2015](#)), or miscarriage rates ( $n = 9$ , RR 1.092, 95% CI 0.846–1.408,  $I^2 = 0\%$ ; [Supplementary Fig. S5d](#); [Mansour et al., 2011](#); [HoNg et al., 2014](#); [Singh and Singh, 2014](#); [Zarei et al., 2014](#); [Aaleysin et al., 2015](#); [Wirleitner et al., 2015a,b](#); [Dehghani Firouzabadi et al., 2016](#); [Hosseini et al., 2016](#); [Hafezi et al., 2018](#)). However, there was a significant increase in biochemical pregnancy rate ( $n = 5$ , RR 1.241, 95% CI 1.012–1.521,  $I^2 = 70.33\%$ ; [Supplementary Fig. S5b](#); [Santibañez et al., 2014](#); [Aaleysin et al., 2015](#); [Wirleitner et al., 2015a](#); [Hafezi et al., 2018](#); [Laokirkkiat et al., 2019](#)). Several hCG doses were used ranging from 100 to 6500 IU ([Supplementary Table S1](#)). However, the most commonly used dose was 500 IU (dissolved in embryo culture medium) evaluated in 15 studies, which showed a consistent increase in



clinical pregnancy rates ( $n=15$ , RR 1.275, 95% CI 1.119, 1.4521,  $I^2=60.05\%$ ; Supplementary Fig. S5f; Mansour *et al.*, 2011; Cambiaghi *et al.*, 2013; Leao *et al.*, 2013; HoNg *et al.*, 2014; Santibañez *et al.*, 2014; Singh and Singh, 2014; Aaleysin *et al.*, 2015; Wirleitner *et al.*, 2015a,b; Dehghani Firouzabadi *et al.*, 2016; Eskandar *et al.*, 2016; Hosseini *et al.*, 2016; Hafezi *et al.*, 2018; Laokirkkiat *et al.*, 2019; WaNg *et al.*, 2019).

### Hyaluronic acid

In total, nine RCTs evaluated the use of hyaluronic acid (HA) at the time of ET compared to no intervention/placebo (Feichtinger *et al.*, 1992; Simon *et al.*, 2003; Friedler *et al.*, 2005, 2007; Korosec *et al.*, 2007; Mahani and Davar, 2007; Tomari *et al.*, 2014; Fasano *et al.*, 2016; Perez *et al.*, 2019). The random-effect meta-analysis showed a significant increase in clinical pregnancy with the administration of HA versus placebo (RR 1.457, 95% CI 1.197–1.261,  $I^2=46.48\%$ ; Fig. 4a; Feichtinger *et al.*, 1992; Simon *et al.*, 2003; Friedler *et al.*, 2005, 2007; Korosec *et al.*, 2007; Mahani and Davar, 2007; Tomari *et al.*, 2014; Fasano *et al.*, 2016; Perez *et al.*, 2019). Compared to no intervention, HA was seen to significantly increase live birth rate ( $n=4$ , RR 1.471, 95% CI 1.092–1.982,  $I^2=4.89\%$ ; Fig. 4b; Supplementary Fig. S6c; Simon *et al.*, 2003; Korosec *et al.*, 2007; Kandari, 2019; Perez *et al.*, 2019), but did not affect ongoing pregnancy rate ( $n=4$ , RR 1.170, 95% CI 0.894–1.532,  $I^2=0.01\%$ ; Supplementary Fig. S6b; Feichtinger *et al.*, 1992; Simon *et al.*, 2003; Friedler *et al.*, 2007; Korosec *et al.*, 2007).

Twelve RCTs compared the use of a high- versus low-dose HA at the time of ET (Schoolcraft *et al.*, 2002; Balaban *et al.*, 2004; Yakin *et al.*, 2004; Ravhon *et al.*, 2005; Walker *et al.*, 2005; Valojerdi *et al.*, 2006; Morbeck, 2007; Urman *et al.*, 2008; Dittmann-Müller *et al.*, 2009; Fancsovits *et al.*, 2011, 2015; Yung *et al.*, 2021), which was associated with a mild increase in clinical pregnancy (RR 1.097, 95% CI 1.028–1.169,  $I^2=0.00\%$ ; Supplementary Fig. S7a; Schoolcraft *et al.*, 2002; Balaban *et al.*, 2004; Yakin *et al.*, 2004; Ravhon *et al.*, 2005; Walker *et al.*, 2005; Valojerdi *et al.*, 2006; Morbeck, 2007; Urman *et al.*, 2008; Dittmann-Müller *et al.*, 2009; Fancsovits *et al.*, 2011, 2015; Yung *et al.*, 2021), but had no significant effect on other reproductive outcomes (ongoing pregnancy rate ( $n=2$ , RR 0.918, 95% CI 0.714–1.179,  $I^2=3.63\%$ ; Morbeck, 2007; Yung *et al.*, 2021) and live birth rate ( $n=6$ , RR 1.134, 95% CI 1.000–1.285,  $I^2=22.17\%$ ; Valojerdi *et al.*, 2006; Morbeck, 2007; Urman *et al.*, 2008; Fancsovits *et al.*, 2011, 2015; Yung *et al.*, 2021; Supplementary Fig. S7b, c)).

### Granulocyte colony-stimulating factor

Our meta-analysis of four RCTs evaluating the effects of using granulocyte colony-stimulating factor (G-CSF) at the time of ET compared to placebo (Singh *et al.*, 2015; Obidniak *et al.*, 2016; Arefi *et al.*, 2018; Singh and Singh, 2018) showed a significant increase in clinical pregnancy (RR 1.774, 95% CI 1.252–2.512,  $I^2=0$ ; Fig. 4a, Supplementary Fig. S8). Only one RCT reported on each of live birth (RR 1.518, 95% CI 0.769–2.997; Arefi *et al.*, 2018), miscarriage (RR 1.272, 95% CI 0.123–13.177; Arefi *et al.*, 2018) and ongoing pregnancy (RR 2.473, 95% CI 1.131–5.405; Singh and Singh, 2018), all showing mixed benefit of using G-CSF with low evidence quality (Supplementary Table SIV).

### Seminal fluid

The use of seminal fluid was assessed in two RCTs (Aflatoonian *et al.*, 2009; Karimian *et al.*, 2010) of medium risk of bias (Supplementary Table SIII). There was no significant benefit to improve biochemical pregnancy ( $n=2$ , RR 1.138, 95% CI 0.912, 1.420,  $I^2=0\%$ ; Supplementary Fig. S9), clinical pregnancy (RR 1.205, 95% CI 0.717–2.025) or live birth (RR 1.281, 95% CI 0.908–1.808; Aflatoonian *et al.*, n.d.; Karimian *et al.*, 2010).

### Culture medium

One RCT (Sigalos *et al.*, 2018) evaluated the impact of ET low versus high volumes of culture medium, which showed no benefit on clinical pregnancy (RR 0.862, 95% CI 0.662–1.125) or ongoing pregnancy (RR 0.833, 95% CI 0.586–1.185).

### Plasma infusion

The study by Obidniak *et al.* (2017) evaluated the effects of plasma infusion at the time of ET ( $n=90$  participants) reporting a 2-fold increase in clinical pregnancy compared to placebo (RR 2.182, 95% CI 1.219–3.904). Confidence in this evidence was judged to be low (Supplementary Table SIII).

### 17-Hydroxyprogesterone caproate

We identified one RCT (Abu-Musa *et al.*, 2008) that evaluated the effectiveness of providing one injection of 17-hydroxyprogesterone caproate (17-HPC) compared to placebo at the time of ET ( $n=125$  participants). There was no significant effect on clinical pregnancy with 17-HPC injection versus placebo (RR 0.902, 95% CI 0.569, 1.429).


### Powdered gloves

A single RCT evaluated the difference in using powdered versus unpowdered gloves at the time of ET ( $n=712$  participants; Hannoun *et al.*, 2009). There was no difference in clinical pregnancy rate across both groups (RR 1.008, 95% CI 0.834–1.218).

## ET techniques

### Use of ultrasound

A large body of evidence was captured evaluating the different ultrasound modalities used at the time of ET. We pooled data from 24 RCTs that compared ET under ultrasound guidance to ET guided by clinical touch, which showed a significant increase in clinical pregnancy ( $n=24$ , RR 1.265, 95% CI 1.151–1.391,  $I^2=8.53\%$ ; Fig. 4a; Wisanto *et al.*, 1989; Prapas *et al.*, 1995; Kan *et al.*, 1999; Coroleu *et al.*, 2000, 2002b; Abdelmassih *et al.*, 2001; Garc'a-Velasco *et al.*, 2001, 2002; Tang *et al.*, 2001; Matorras *et al.*, 2002; Sallam *et al.*, 2002; Bar-Hava *et al.*, 2003; Marconi *et al.*, 2003; Weissman *et al.*, 2003; de Camargo Martins *et al.*, 2004; Moraga-Sanchez *et al.*, 2004; Li *et al.*, 2005; Maldonado *et al.*, 2005; Chen *et al.*, 2007; Davar *et al.*, 2007; Kosmas *et al.*, 2007; Drakeley *et al.*, 2008; Eskandar *et al.*, 2008; Ammar *et al.*, 2013), ongoing pregnancy ( $n=8$ , RR 1.369, 95% CI 1.144–1.639,  $I^2=51.34\%$  (Supplementary Fig. S10c; Coroleu *et al.*, 2000; Tang *et al.*, 2001; Matorras *et al.*, 2002; Marconi *et al.*, 2003; Davar *et al.*, 2007; Drakeley *et al.*, 2008; Eskandar *et al.*, 2008; Ammar *et al.*, 2013), biochemical pregnancy ( $n=2$ , RR 1.496, 95% CI 1.174–1.906; Supplementary Fig. S10b; Coroleu *et al.*, 2000; Kosmas *et al.*, 2007), and live birth ( $n=2$ , RR 1.744, 95% CI 1.163–2.706,  $I^2=79.97\%$ ; Fig. 4b; Supplementary Fig. S10e; Eskandar *et al.*, 2008; Azmy *et al.*,



**Figure 4.** Summary forest plots of effect estimates of evaluated interventions at the time of embryo transfer on the reproductive outcomes in women undergoing assisted reproduction. G-CSF, granulocyte colony-stimulating factor; TA, transabdominal; TV, transvaginal; CT, computerized tomography; NSAID, non-steroidal anti-inflammatory drug; RCT, randomized controlled trial; HA, hyaluronic acid.

2009) with the use of ultrasound. There was no difference in miscarriage rates between both groups ( $n=12$ , RR 1.145, 95% CI 0.883–1.484,  $I^2=0\%$ ; [Supplementary Fig. S10d](#); [García-Velasco et al., 2001, 2002](#); [Tang et al., 2001](#); [Coroleu et al., 2002b](#); [Matorras et al., 2002](#); [Weissman et al., 2003](#); [de Camargo Martins et al., 2004](#); [Davar et al., 2007](#); [Kosmas et al., 2007](#); [Eskandar et al., 2008](#); [Azmy et al., 2009](#); [Ammar et al., 2013](#)).

We included seven RCTs that compared the use of transabdominal versus transvaginal ultrasound at the time of ET ([Porat et al., 2010](#); [Bodri et al., 2011](#); [Deep et al., 2013](#); [Hauzman et al., 2013](#); [Dalal et al., 2014](#); [Revelli et al., 2016](#); [Karavani et al., 2017](#)). There was no difference between both groups on clinical pregnancy ( $n=7$ , RR

1.004, 95% CI 0.924–1.090,  $I^2=0\%$ ; [Supplementary Fig. S10f](#); [Porat et al., 2010](#); [Bodri et al., 2011](#); [Deep et al., 2013](#); [Hauzman et al., 2013](#); [Dalal et al., 2014](#); [Revelli et al., 2016](#); [Karavani et al., 2017](#)), ongoing pregnancy ( $n=5$ , RR 1.029, 95% CI 0.922–1.148,  $I^2=0.01\%$ ; [Supplementary Fig. S10h](#); [Porat et al., 2010](#); [Bodri et al., 2011](#); [Hauzman et al., 2013](#); [Revelli et al., 2016](#); [Karavani et al., 2017](#)), biochemical pregnancy ( $n=2$ , RR 0.977, 95% CI 0.712–1.341; [Supplementary Fig. S10g](#); [Bodri et al., 2011](#); [Hauzman et al., 2013](#)), miscarriage ( $n=3$ , RR 0.835, 95% CI 0.575–1.211,  $I^2=0\%$ ; [Supplementary Fig. S10i](#); [Bodri et al., 2011](#); [Revelli et al., 2016](#); [Karavani et al., 2017](#)) and live birth ( $n=1$ , RR 0.789, 95% CI 0.444–1.403; [Karavani et al., 2017](#)). One RCT by [Saravolos et al. \(2016\)](#)

evaluated the use of 3-dimensional versus 2-dimensional ultrasound at the time of ET ( $n = 474$  participants), which showed no difference for clinical pregnancy (RR 0.981, 95% CI 0.800–1.202) or any of the other secondary outcomes.

### Bladder fullness

Three RCTs compared the difference in reproductive outcomes following ET on a full versus empty bladder ( $n = 1069$  participants; Mitchell *et al.*, 1989; Lewin *et al.*, 1997; Lorusso *et al.*, 2005). Overall, there was no difference across both groups for clinical pregnancy ( $n = 3$ , RR 1.266, 95% CI 0.884–1.813,  $I^2 = 50.68\%$ ; Supplementary Fig. S11a; Mitchell *et al.*, 1989; Lewin *et al.*, 1997; Lorusso *et al.*, 2005), ongoing pregnancy ( $n = 2$ , RR 1.276, 95% CI 0.874–1.863; Supplementary Fig. S11b; Lewin *et al.*, 1997; Lorusso *et al.*, 2005) or miscarriage ( $n = 1$ , RR 1.047, 95% CI 0.467–2.346; Lorusso *et al.*, 2005).

### Pressure on cervix

Two RCTs evaluated the effect of mechanical effect on the cervix by unscrewing the speculum to apply gentle pressure during ET ( $n = 716$  participants), which showed no significant effect for clinical pregnancy ( $n = 2$ , RR 1.175, 95% CI 0.817–1.690; Supplementary Fig. S12; Mansour, 2005; Amui *et al.*, 2011) or ongoing pregnancy ( $n = 1$ , RR 0.843, 95% CI 0.485–1.468; Amui *et al.*, 2011).

### Pump regulated transfer

One study (Caanen *et al.*, 2016) compared the use of a pump regulated transfer to manual ET ( $n = 599$  participants), which showed no difference between methods for clinical pregnancy (RR 1.201, 95% CI 0.904–1.596) or ongoing pregnancy (RR 1.234, 95% CI 0.884–1.722).

### Cervical mucus removal

Removing the cervical mucus before ET was assessed in six studies (Ruhlman *et al.*, 1999; Soroka *et al.*, 1999; Glass *et al.*, 2000; Berkkanoglu *et al.*, 2006; Visschers *et al.*, 2007; Moini *et al.*, 2011) compared to standard practice ( $n = 1157$  participants) using different tools including cotton swabs (Moini *et al.*, 2011), cervical brush (Visschers *et al.*, 2007), aspirators (Ruhlman *et al.*, 1999; Soroka *et al.*, 1999) or flushing with culture media solution (Berkkanoglu *et al.*, 2006) or their combination (Glass *et al.*, 2000). Our meta-analysis showed no significant effect of cervical mucus removal on clinical pregnancy ( $n = 5$ , RR 1.029, 95% CI 0.753–1.405,  $I^2 = 5.89\%$ ; Supplementary Fig. S13a; Ruhlman *et al.*, 1999; Soroka *et al.*, 1999; Glass *et al.*, 2000; Berkkanoglu *et al.*, 2006; Moini *et al.*, 2011), biochemical pregnancy ( $n = 2$ , RR 0.933, 95% CI 0.774–1.125; Supplementary Fig. S13b; Visschers *et al.*, 2007; Moini *et al.*, 2011), ongoing pregnancy ( $n = 2$ , RR 1.068, 95% CI 0.849–1.344; Supplementary Fig. S13c; Berkkanoglu *et al.*, 2006; Visschers *et al.*, 2007), miscarriage ( $n = 2$ , RR 0.790, 95% CI 0.487–1.282; Supplementary Fig. S13d; Visschers *et al.*, 2007; Moini *et al.*, 2011) or live birth ( $n = 2$ , RR 1.506, 95% CI 0.908–2.499; Supplementary Fig. S13e; Visschers *et al.*, 2007; Moini *et al.*, 2011).

### ET catheters

A total of 27 RCTs compared ET outcomes using soft versus hard catheters (Gilberto Almodin *et al.*, n.d.; Wisanto *et al.*, 1989; Al-Shawaf *et al.*, 1993; Perin *et al.*, 1997; Grunert *et al.*, 1998; Amorcho *et al.*, 1999; Ghazzawi *et al.*, 1999; Mayer *et al.*, 1999; Boone *et al.*,

2001; Curfs *et al.*, 2001; Lavery *et al.*, 2001; Karande *et al.*, 2002; Levi-Setti *et al.*, 2002; McDonald and Norman, 2002; Mortimer *et al.*, 2002; van Weering *et al.*, 2002; Foutouh *et al.*, 2003; McIlveen *et al.*, 2005; Baris *et al.*, 2007; Rhodes *et al.*, 2007; El-Shawarby *et al.*, 2008; Saldeen *et al.*, 2008; Yao *et al.*, 2009b; Allahbadia *et al.*, 2010; Talwar *et al.*, 2011; Nihat Candan *et al.*, 2014). Overall, there was a significant increase in clinical pregnancy with the use of a soft catheter (RR 1.122, 95% CI 1.028–1.224,  $I^2 = 57.66\%$ ; Fig. 4a; Supplementary Fig. S14a) but no difference was found for ongoing pregnancy ( $n = 3$ , RR 1.138, 95% CI 0.904–1.432,  $I^2 = 32.46\%$ ; Supplementary Fig. S14b; Karande *et al.*, 2002; Levi-Setti *et al.*, 2002; Baris *et al.*, 2007) or live birth rates ( $n = 2$ , RR 2.222, 95% CI 0.457–10.806,  $I^2 = 94.13\%$ ; Fig. 4b; Supplementary Fig. S14c; Perin *et al.*, 1997; Saldeen *et al.*, 2008).

Using a double compared to a single lumen ET catheter seemed to reduce clinical pregnancy rate in a small RCT of 66 participants (RR 0.314, 95% CI 0.129–0.764; Meriano *et al.*, 2000). Similarly, using a straight versus a bent tip catheter seemed to also reduce clinical pregnancy in another small RCT (Ocal *et al.*, 2003; RR 0.648, 95% CI 0.421–0.999). The use of an echogenic versus standard ET catheter did not affect the rate of clinical pregnancy in both groups (RR 1.302, 95% CI 0.966–1.756; Coroleu *et al.*, 2006). Confidence in evidence sought on these interventions was low (Table 1, Supplementary Table SIV). Yayla Abide *et al.* (2018) suggested that catheter rotation at the time of ET compared to standard practice yielded a higher clinical pregnancy rate (RR 1.560, 95% CI 1.026–2.371) with no difference reported for ongoing pregnancy (RR 1.435, 95% CI 0.911–2.260) or biochemical pregnancy (RR 1.577, 95% CI 1.051–2.366). This RCT showed a moderate risk of bias and its results should be interpreted with caution (Supplementary Table SIII).

Three RCTs compared the use of air or fluid in the transfer catheter (Krampl *et al.*, 1995; Moreno *et al.*, 2004; Madani *et al.*, 2010). Both interventions had a similar effect on clinical pregnancy rate (RR 1.361, 95% CI 0.844–2.195) with moderate inconsistency ( $I^2 = 61.16\%$ ; Supplementary Fig. S15). We included three RCTs that compared early versus delayed catheter withdrawal after ET (Martínez *et al.*, 2001; Arvas *et al.*, 2014; Devranoğlu *et al.*, 2018). Both techniques yielded a similar clinical pregnancy rate ( $n = 3$ , RR 1.039, 95% CI 0.868–1.243,  $I^2 = 0\%$ ; Supplementary Fig. S16; Martínez *et al.*, 2001; Arvas *et al.*, 2014; Devranoğlu *et al.*, 2018), ongoing pregnancy ( $n = 1$ , RR 0.889, 95% CI 0.594–1.329; Devranoğlu *et al.*, 2018) and biochemical pregnancy ( $n = 1$ , RR 1.014, 95% CI 0.728–1.413; Devranoğlu *et al.*, 2018).

The site of embryo deposition ( $>15$  mm versus  $<15$  mm from the fundus) was assessed in three RCTs (Nazari *et al.*, 1993; Coroleu *et al.*, 2002a; Franco *et al.*, 2004). Our pooled effect estimate showed no difference between both groups for clinical pregnancy ( $n = 3$ , RR 1.089, 95% CI 0.828–1.431,  $I^2 = 60.99\%$ ; Supplementary Fig. S17a; Nazari *et al.*, 1993; Coroleu *et al.*, 2002a; Franco *et al.*, 2004), ongoing pregnancy ( $n = 3$ , RR 1.191, 95% CI 0.858–1.653,  $I^2 = 65.08\%$ ; Supplementary Fig. S17b; Nazari *et al.*, 1993; Coroleu *et al.*, 2002a; Franco *et al.*, 2004), miscarriage ( $n = 2$ , RR 0.916, 95% CI 0.446–1.881; Supplementary Fig. S17c; Coroleu *et al.*, 2002a; Franco *et al.*, 2004) or live birth ( $n = 3$ , RR 1.201, 95% CI 0.855–1.687,  $I^2 = 66.36\%$ ; Supplementary Fig. S17d; Nazari *et al.*, 1993; Coroleu *et al.*, 2002a; Franco *et al.*, 2004). One small RCT (Groutz *et al.*, 1997;  $n = 40$  participants) compared the outcome of transmyometrial



**Table 1** GRADE assessment of evidence from randomised trials evaluating interventions at the time of embryo transfer in women undergoing assisted reproduction

Certainty assessment					Summary of findings			
Participants (studies) Follow up	Risk of bias	Inconsistency	Indirectness	Imprecision	Overall certainty of evidence	Study event rates (%)		Relative effect (95% CI)
						With Comparison	With Interventions at the time of ET	
Higher confidence and significant effect size:								
Hyaluronic acid versus no HA								
2453 (9 RCTs)	not serious	not serious	not serious	not serious	⊕⊕⊕⊕ HIGH	286/1305 (21.9%)	387/1148 (33.7%)	RR 1.457 (1.197 to 1.773)
Ultrasound guided ET versus clinical touch								
7256 (24 RCTs)	not serious	not serious	not serious	serious <sup>a</sup>	⊕⊕⊕○ MODERATE	986/3587 (27.5%)	1245/3669 (33.9%)	RR 1.265 (1.151 to 1.391)
Softer versus harder catheters								
10688 (27 RCTs)	not serious	not serious	serious <sup>b</sup>	not serious	⊕⊕⊕○ MODERATE	1704/5248 (32.5%)	1969/5440 (36.2%)	RR 1.122 (1.028 to 1.224)
Lower confidence and significant effect size:								
G-CSF versus placebo/ no G-CSF								
292 (4 RCTs)	serious <sup>c</sup>	not serious	not serious	serious <sup>a</sup>	⊕⊕○○ LOW	35/154 (22.7%)	58/138 (42.0%)	RR 1.774 (1.252 to 2.512)
Atosiban versus placebo/no atosiban								
1646 (7 RCTs)	not serious	serious <sup>d</sup>	not serious	serious <sup>a</sup>	⊕⊕○○ LOW	325/823 (39.5%)	426/823 (51.8%)	RR 1.493 (1.184 to 1.882)
hCG versus placebo/no hCG								
4653 (16 RCTs)	serious <sup>e</sup>	serious <sup>f</sup>	not serious	not serious	⊕⊕○○ LOW	883/2298 (38.4%)	1052/2355 (44.7%)	RR 1.232 (1.099 to 1.382)
High confidence and no significant effect:								
Transvaginal ultrasound versus transabdominal ultrasound								
2942 (7 RCTs)	not serious	not serious	not serious	not serious	⊕⊕⊕⊕ HIGH	620/1457 (42.6%)	638/1485 (43.0%)	RR 1.004 (0.924 to 1.090)
Some positive associations with clinical pregnancy rate:								
NSAID's versus placebo/ no NSAID's								
1207 (7 RCTs)	not serious	not serious	serious <sup>h</sup>	not serious	⊕⊕⊕○ MODERATE	166/599 (27.7%)	222/608 (36.5%)	RR 1.294 (0.973 to 1.721)
Continued								

Continued

Table I Continued

Certainty assessment					Summary of findings			
Participants (studies) Follow up	Risk of bias	Inconsistency	Indirectness	Imprecision	Overall certainty of evidence	Study event rates (%)		Relative effect (95% CI)
						With Comparison	With Interventions at the time of ET	
Acupuncture versus placebo/no acupuncture								
6496 (18 RCTs)	not serious	serious <sup>i</sup>	not serious	not serious	⊕⊕⊕⊕ MODERATE	1379/3409 (40.5%)	1379/3087 (44.7%)	<b>RR 1.121</b> (0.988 to 1.273)
Air versus fluid in catheter tip								
408 (3 RCTs)	serious <sup>c</sup>	serious <sup>j</sup>	not serious	serious <sup>a</sup>	⊕⊕⊕⊕ VERY LOW	58/203 (28.6%)	76/205 (37.1%)	<b>RR 1.361</b> (0.844 to 2.195)
Full bladder versus empty bladder								
1069 (3 RCTs)	serious <sup>c</sup>	serious <sup>k</sup>	not serious	very serious <sup>g</sup>	⊕⊕⊕⊕ VERY LOW	102/525 (19.4%)	148/544 (27.2%)	<b>RR 1.266</b> (0.884 to 1.813)
Lower confidence and no real associations with clinical pregnancy rate:								
Music versus no music								
409 (3 RCTs)	not serious	not serious	not serious	very serious <sup>g</sup>	⊕⊕⊕⊕ LOW	101/209 (48.3%)	102/200 (51.0%)	<b>RR 1.052</b> (0.866 to 1.279)
Nurse versus doctor								
655 (2 RCTs)	not serious	not serious	not serious	very serious <sup>g</sup>	⊕⊕⊕⊕ LOW	115/327 (35.2%)	111/328 (33.8%)	<b>RR 0.961</b> (0.778 to 1.187)
Pressure on cervix versus no pressure								
716 (2 RCTs)	serious <sup>c</sup>	not serious	not serious	very serious <sup>g</sup>	⊕⊕⊕⊕ VERY LOW	172/359 (47.9%)	221/357 (61.9%)	<b>RR 1.175</b> (0.817 to 1.690)
Site of deposition: further from fundus (15–20mm) versus closer (<12mm)								
2170 (3 RCTs)	serious <sup>c</sup>	serious <sup>l</sup>	very serious <sup>m</sup>	very serious <sup>g</sup>	⊕⊕⊕⊕ VERY LOW	176/921 (19.1%)	256/1249 (20.5%)	<b>RR 1.089</b> (0.828 to 1.431)
Delayed versus immediate catheter withdrawal								
672 (3 RCTs)	serious <sup>c</sup>	not serious	not serious	very serious <sup>g</sup>	⊕⊕⊕⊕ VERY LOW	129/339 (38.1%)	128/333 (38.4%)	<b>RR 1.039</b> (0.868 to 1.243)
Cervical mucus removal versus no removal								
1157 (5 RCTs)	serious <sup>c</sup>	serious <sup>o</sup>	not serious	very serious <sup>g</sup>	⊕⊕⊕⊕ VERY LOW	225/597 (37.7%)	229/560 (40.9%)	<b>RR 1.029</b> (0.753 to 1.405)
Antibiotics versus no antibiotics								
625 (2 RCTs)	serious <sup>c</sup>	not serious	serious <sup>p</sup>	very serious <sup>g</sup>	⊕⊕⊕⊕ VERY LOW	109/317 (34.4%)	107/308 (34.7%)	<b>RR 1.008</b> (0.812 to 1.251)

Continued

Table 1 Continued

Participants (studies) Follow up	Certainty assessment				Summary of findings		
	Risk of bias	Inconsistency	Indirectness	Imprecision	Overall certainty of evidence	Study event rates (%)	
						With Comparison	With Interventions at the time of ET
<b>Mindfulness versus no mindfulness</b>							
500 (2 RCTs)	serious <sup>c</sup>	not serious	not serious	very serious <sup>g</sup>	⊕○○○ VERY LOW	121/247 (49.0%)	113/253 (44.7%)  <b>RR 0.911</b> (0.755 to 1.098)
<b>Significant negative association with clinical pregnancy rates:</b>							
<b>Bed rest versus shorter/no bed rest</b>							
1180 (6 RCTs)	serious <sup>c</sup>	not serious	not serious	serious <sup>a</sup>	⊕⊕○○ LOW	217/591 (36.7%)	185/589 (31.4%)  <b>RR 0.857</b> (0.741 to 0.991)

The table presents evidence summary for interventions evaluated in two or more randomised trials.

GRADE: Grading of Recommendations Assessment, Development and Evaluation, CI: Confidence interval, RR: Risk ratio, ET: embryo transfer, G-CSF: granulocyte colony-stimulating factor, HA: hyaluronic acid, NSAID: non-steroidal anti-inflammatory drug

**Explanations**

<sup>a</sup>Wide confidence intervals

<sup>b</sup>Significant heterogeneity in study design: lots of different catheter types compared which we believe may have impacted outcomes

<sup>c</sup>Downgraded due to concerns about whole risk of bias

<sup>d</sup>Downgraded due to high heterogeneity: I<sup>2</sup> = 68.27%

<sup>e</sup>Downgraded due to concerns about allocation concealment and lack of blinding

<sup>f</sup>Downgraded due to high heterogeneity: I<sup>2</sup> = 57.76%

<sup>g</sup>Very wide confidence intervals

<sup>h</sup>Significant heterogeneity in study design: lots of different NSAID's and length of treatment

<sup>i</sup>Downgraded due to high heterogeneity: I<sup>2</sup> = 75.14%

<sup>j</sup>Downgraded due to high heterogeneity: I<sup>2</sup> = 61.16%

<sup>k</sup>Downgraded due to high heterogeneity: I<sup>2</sup> = 50.68%

<sup>l</sup>Downgraded due to high heterogeneity: I<sup>2</sup> = 60.99%

<sup>m</sup>Significant heterogeneity in study design, not consistent site of deposition in any of them

<sup>n</sup>Downgraded due to high heterogeneity: I<sup>2</sup> = 75.89%

to transcervical ultrasound-guided ET showing no difference in the rate of clinical pregnancy (RR 0.333, 95% CI 0.038–2.939) or ongoing pregnancy (RR 0.143, 95% CI 0.008–2.599).

### Bed rest

We performed a meta-analysis of six RCTs that evaluated the effect of bed rest following ET compared to immediate ambulation or no rest (Botta and Grudzinskas, 1997; Rezábek et al., 2001; Amarín and Obeidat, 2004; Purcell et al., 2007; Gaikwad et al., 2013; Malhotra and Sarkar, 2019). Overall, there was a reduction in clinical pregnancy rates in the bed rest group (RR 0.857, 95% CI 0.741–0.991,  $I^2 = 0.01\%$ ; Fig. 4a, Supplementary Fig. S18a) while there was no significant difference for biochemical pregnancy ( $n = 2$ , RR 0.823, 95% CI 0.538–1.259; Supplementary Fig. S18b; Amarín and Obeidat, 2004; Gaikwad et al., 2013), ongoing pregnancy ( $n = 4$ , RR 0.870, 95% CI 0.656–1.153,  $I^2 = 50.53\%$ ; Supplementary Fig. S18c; Botta and Grudzinskas, 1997; Amarín and Obeidat, 2004; Purcell et al., 2007; Gaikwad et al., 2013), miscarriage ( $n = 3$ , RR 1.076, 95% CI 0.466–2.483,  $I^2 = 61.84\%$ ; Supplementary Fig. S18d; Botta and Grudzinskas, 1997; Amarín and Obeidat, 2004; Gaikwad et al., 2013) or live birth ( $n = 3$ , RR 0.800, 95% CI 0.626–1.022,  $I^2 = 21.93\%$ ; Fig. 4b; Supplementary Fig. S18e; Rezábek et al., 2001; Gaikwad et al., 2013; Malhotra and Sarkar, 2019).

### Nurse versus doctor

Two RCTs compared ET outcomes when performed by a nurse or a doctor ( $n = 655$  participants) showing no significant difference between both groups for clinical pregnancy ( $n = 2$ , RR of 0.961, 95% CI 0.778–1.187; Supplementary Fig. S19; Bjuresten et al., 2003; Rinaldi et al., 2014) or ongoing pregnancy ( $n = 1$ , RR 0.981, 95% CI 0.728–1.324; Rinaldi et al., 2014).

## Relaxation interventions

### Acupuncture

Eighteen RCTs evaluated the effect of acupuncture interventions on clinical pregnancy ( $n = 6437$  participants) including 16 studies on needle acupuncture (Paulus et al., 2002, 2003; Benson et al., 2006; Westergaard et al., 2006; Craig et al., 2007, 2014; Fratterelli et al., 2008; Domar et al., 2009; So et al., 2009, 2010; Andersen et al., 2010; Madaschi et al., 2010; Omodei et al., 2010; Moy et al., 2011; Qu et al., 2014; Dehghani et al., 2020), two on laser acupuncture (Benson et al., 2006; Fratterelli et al., 2008) and two on transcutaneous electrical acupoint stimulation (TEAS; Zhang et al., 2011; Zhong and Zhang, 2017). Eight studies compared acupuncture to sham acupuncture therapy (Paulus et al., 2003; Benson et al., 2006; Fratterelli et al., 2008; So et al., 2009, 2010; Andersen et al., 2010; Moy et al., 2011; Zhang et al., 2011), whilst 12 of the studies compared acupuncture to no treatment (Paulus et al., 2002; Benson et al., 2006; Westergaard et al., 2006; Craig et al., 2007, 2014; Fratterelli et al., 2008; Domar et al., 2009; Madaschi et al., 2010; Omodei et al., 2010; Qu et al., 2014; Zhong and Zhang, 2017; Dehghani et al., 2020). Overall, acupuncture interventions did not increase the rate of clinical pregnancy (RR 1.121, 95% CI 0.988–1.273,  $I^2 = 75.14\%$ ; Supplementary Fig. S20a). Subgroup analysis showed a similar effect for needle ( $n = 16$ , RR 1.076, 95% CI

0.919–1.260,  $I^2 = 73.44\%$ ) and laser acupuncture ( $n = 2$ , RR 1.193, 95% CI 0.996–1.430,  $I^2 = 0\%$ ) whilst TEAS showed some improvement in clinical pregnancy rate ( $n = 2$ , RR 1.345, 95% CI 1.039–1.741,  $I^2 = 62.66\%$ ) but confidence in this evidence was low (Supplementary Fig. S20f).

Compared to standard care or sham intervention, acupuncture showed some benefit in increasing the rate of biochemical pregnancy ( $n = 8$ , RR 1.150, 95% CI 1.002–1.321,  $I^2 = 70.45\%$ ; Benson et al., 2006; Westergaard et al., 2006; Craig et al., 2007; Fratterelli et al., 2008; Domar et al., 2009; Zhang et al., 2011; Zhong and Zhang, 2017; Dehghani et al., 2020) though evidence was of poor quality with high heterogeneity. There was no difference for ongoing pregnancy ( $n = 11$ , RR 1.130, 95% CI 0.931–1.371,  $I^2 = 75.16\%$ ; Paulus et al., 2002, 2003; Westergaard et al., 2006; Fratterelli et al., 2008; So et al., 2009, 2010; Andersen et al., 2010; Madaschi et al., 2010; Omodei et al., 2010; Seto et al., 2017; Dehghani et al., 2020), miscarriage ( $n = 10$ , RR 1.073, 95% CI 0.828–1.391,  $I^2 = 0.0\%$ ; Paulus et al., 2002, 2003; Westergaard et al., 2006; So et al., 2009, 2010; Andersen et al., 2010; Madaschi et al., 2010; Omodei et al., 2010; Zhang et al., 2011; Craig et al., 2014) or live birth ( $n = 11$ , RR 1.176, 95% CI 0.929–1.488,  $I^2 = 80.92\%$ ; Paulus et al., 2002, 2003; So et al., 2009, 2010; Andersen et al., 2010; Madaschi et al., 2010; Omodei et al., 2010; Zhang et al., 2011; Craig et al., 2014; Qu et al., 2014; Seto et al., 2017; Fig. 4b, Supplementary Fig. S20b–e). There were no significant differences in clinical pregnancy rates observed between studies using a sham acupuncture control ( $n = 8$ , RR 1.056, 95% CI 0.890–1.252); or those with no placebo comparison ( $n = 12$ , RR 1.177, 95% CI 0.988–1.273; Supplementary Fig. S20g).

Listening to music during ET was assessed in three RCTs compared to standard care (Murphy et al., 2014; Stocker et al., 2016; Aba et al., 2017). Overall, this did not increase the rate of clinical pregnancy (RR 1.052, 95% CI 0.866–1.279,  $I^2 = 0.00\%$ ; Supplementary Fig. S21). Similarly, the use of mindfulness was assessed in two RCTs (Benson et al., 2006; Fratterelli et al., 2008) also showing no difference in clinical pregnancy (RR 0.911, 95% CI 0.755–1.098,  $I^2 = 0.00\%$ ) or biochemical pregnancy rate (RR 0.912, 95% CI 0.780–1.066,  $I^2 = 0\%$ ; Supplementary Fig. S22a and b, respectively). The effect of mindfulness on ongoing pregnancy rate was reported in one study (Fratterelli et al., 2008) showing no significant increase was seen (RR 0.946, 95% CI 0.738–1.211).

One study measured the effect of porcine relaxin compared to a placebo (MacLennan et al., 1985). There was no increase in clinical pregnancy (RR 0.882, 95% CI 0.405–1.924) or ongoing pregnancy rate (RR 1.177, 95% CI 0.442–3.134) or miscarriage rate (RR 0.441, 95% CI 0.085–2.296). Similarly, one study measured the effect of glyceryl trinitrate compared to a placebo (Shaker et al., 1993) also showing no increase in clinical pregnancy rate (RR 0.947, 95% CI 0.554–1.620).

There was one study that investigated massage therapy as an intervention compared to standard care (Gavrizi and Skilleren, 2019). The results from this study showed no significant increase in live birth with massage (RR 1.371, 95% CI 0.881–2.133). Catoire et al. (2013) compared the effect of hypnosis on the day of ET to the use of benzodiazepine in 93 women, which did not report a significant difference for clinical pregnancy (RR 1.032, 95% CI 0.595–1.790) or live birth (RR 1.015, 95% CI 0.509–2.03).

## Discussion

### Summary of main findings

In this comprehensive systematic review, we identified a large number of competing interventions proposed to increase the chances of conception following ET. Overall, the quality of evidence was low with significant concerns regarding trial methodology and risk of publication bias. Several interventions do seem, however, to significantly improve the chances of conception (Table I).

The use of ultrasound guidance and soft ET catheters are commonly used in clinical practice with good quality evidence to support their effectiveness. We identified several promising pharmacological agents that also seem to aid embryo implantation and increase clinical pregnancy including HA, G-CSF, atosiban and hCG. A common mechanistic effect across these interventions could be through maximizing uterine relaxation at the time of ET and optimizing endometrial receptivity (Mains and van Voorhis, 2010). Evidence for these interventions largely stems from small trials and further evaluation in larger trials is warranted before adopting them into routine clinical practice.

Recommending bed rest post-ET was the only intervention in this review that significantly reduced clinical pregnancy rates. This is consistent with ASRM Guidance, which recommends against this intervention (Penzias et al., 2017a). We identified a substantial number of unnecessary interventions that lack sufficient evidence in support of their effectiveness and safety. Such interventions should not be routinely recommended in clinical practice pending assessment in future randomized trials.

### Strengths and limitations

The strengths of our review stem from its pragmatic design and comprehensive search strategy, therefore, offering the most up to date evidence on ET techniques and interventions. We employed a standard methodology and assessed the risk of bias in included trials following a prospectively registered protocol. We conducted subgroup analyses to explore potential effect modifiers, assessed publication bias and employed the GRADE approach to aid evidence translation into clinical practice.

Our findings have several methodological limitations. The quality of included trials was often suboptimal with almost half showing a high risk of bias for outcome reporting (44.9%). This limited our ability to synthesize evidence on all outcomes of interest, particularly live birth. We addressed this perceived bias by downgrading confidence in available evidence in our GRADE assessment of the evaluated interventions; however, this remains a subjective judgement of evidence quality and certainty.

Several effect modifiers could impact the outcome of ET including the operator's experience, patient characteristics (age, BMI, uterine anomalies, etc.), number of embryos transferred and the transfer of fresh versus frozen embryos. Clearly, the technology and practice of assisted conception have also progressed significantly over the time span of included trials (1989–2021) leading to inherent variations in evaluated ART protocols (Supplementary Table SI). We were unable to explore certain effect modifiers due to reporting limitations, however, we attempted to address this by implementing a random-effect model to adjust our pooled effect estimate (Cornell et al., 2014). Further evaluation using individual patient data (IPD) is, therefore, warranted to synthesize higher-quality evidence.

### Implications for clinical practice

Currently, a large number of add-on treatments are proposed to couples undergoing ART, yet the debate on their effectiveness and safety prevail (Zemyarska, 2019; Lensen et al., 2021). We detected many RCTs conducted within the last 20 years evaluating many different interventions focused on optimizing ET success. This highlights the important role of robust evidence synthesis to inform clinical practice and minimize the risk of exposing patients to potentially harmful or unnecessary interventions.

Most of the published systematic reviews on ET techniques are clustered around interventions commonly used in current practice such as ultrasound guidance (Teixeira et al., 2015; Cozzolino et al., 2018) and acupuncture (El-Toukhy et al., 2008; Gu et al., 2019; Smith et al., 2019). While helpful, systematic reviews focused on evaluating singular interventions can lead to fragmented evidence synthesis and an underappreciation of the body of evidence relevant to a particular health condition. To that end, we implemented a comprehensive search strategy, which helped us to detect several interventions of uncertain benefit that were evaluated in one or two RCTs only. We, therefore, caution against the routine uptake of such 'experimental' interventions into clinical practice pending further evaluation in larger, well-conducted RCTs. Although we detected several ET interventions that appear to improve chances of conception, the current body of evidence remains imprecise, and these interventions should be offered in research settings only.

Publication bias and selective outcomes reporting are major barriers to robust evidence synthesis (Sutton et al., 2000) specifically in the context of high stakes interventions such as ET. We chose to report primarily on clinical pregnancy rate to offer the most comprehensive evidence synthesis as most of the included trials did not report on key additional outcomes such as live birth. Bed rest is an important example where varied outcomes reported in primary RCTs (Botta and Grudzinskas, 1997; Rezábek et al., 2001; Amarín and Obeidat, 2004; Purcell et al., 2007; Gaikwad et al., 2013; Malhotra and Sarkar, 2019) could have contributed to the delay in identifying its adverse impact on clinical pregnancy rate following ET. As such, robust evaluation of the confidence in available evidence is crucial before introducing changes to clinical practice (Pottie et al., 2012).

Professional societies should champion the concurrent evaluation of available evidence to regularly update clinical practice guidelines. For example, the most up to date ASRM guidelines on ET recommend cervical mucus removal based on 'fair' evidence at the time (one RCT and one cohort study; Penzias et al., 2017a). Still, our findings summarizing evidence from five RCTs showed no benefit of this intervention.

Finally, the process of ET involves several steps immediately before, during and after where the operator technique seems to significantly impact the final outcome (Cirillo et al., 2020). Establishing a standardized protocol of best practice may facilitate training and reduce variation in practice across fertility clinics (Penzias et al., 2017b).

### Future research need

The majority of the included RCTs in our review had a relatively small sample size with a moderate risk of bias specifically for outcome reporting. This significantly hinders efficient evidence synthesis and increases the risk of small study effect (Harbord et al., 2009). While several of the evaluated interventions seem to have a promising effect in improving clinical pregnancy rate (e.g. Atosiban and hCG), there is a



need to evaluate their effectiveness and safety in well-powered RCTs and prospective IPD meta-analyses to inform clinical practice at scale (Tierney *et al.*, 2015).

In this review, we focused on interventions within 24 h of the ET to increase homogeneity across pooled RCTs. However, several interventions aimed to improve endometrial receptivity and the implantation window could improve the outcome ET outcome such as endometrial scratching and different methods of luteal support (Mains and van Voorhis, 2010). Further evidence synthesis is, therefore, required to advise on the best practice in preparation for ET.

Finally, the majority of the included RCTs did not engage lay consumers in their design, conduct or reporting. Patient engagement is particularly relevant when evaluating anti-anxiety and relaxation interventions during ART treatments, which to date seem to have been based on anecdotal practice (Mahlstedt *et al.*, 1987). Future trials should aim to engage all relevant stakeholders in the ART process to improve the permeation of research into clinical practice.

## Conclusion

Using ultrasound guidance, soft catheters and HA at the time of ET appears to increase pregnancy rates. The use of atosiban, G-CSF and hCG showed a trend towards increased clinical pregnancy rate, but larger trials are required before adopting these interventions in clinical practice. Bed rest post-ET was associated with a reduction in clinical pregnancy and should not be recommended.

## Supplementary data

Supplementary data are available at *Human Reproduction Update* online.

## Data availability

Some or all datasets generated during and/or analysed during the current study are not publicly available but are available from the corresponding author on reasonable request.

## Authors' roles

B.T., H.W. and J.T. conducted the data extraction, primary analysis, data illustration and drafted the first manuscript. S.D.K., D.M. and E.Y. supervised the project conduct and provided critical input to the final manuscript. B.H.A. conceived the idea, wrote the protocol, supervised the analysis and edited the final manuscript.

## Funding

No funding was received towards this work directly. B.H.A. holds a personal development award from the National Institute of Health Research.

## Conflict of interest

Nothing to disclose.

## References

- Aaleya A, Aghahosseini M, Rashidi M, Safdarian L, Sarvi F, Najmi Z, Mobasser A, Amoozgar B. In vitro fertilization outcome following embryo transfer with or without preinstillation of human chorionic gonadotropin into the uterine cavity: a randomized controlled trial. *Gynecol Obstet Invest* 2015;**79**:201–205.
- Aba YA, Avci D, Guzel Y, Ozcelik SK, Gurtekin B. Effect of music therapy on the anxiety levels and pregnancy rate of women undergoing in vitro fertilization-embryo transfer: a randomized controlled trial. *Appl Nurs Res* 2017;**36**:19–24.
- Abdelmassih VG, Abdelmassih ST, Nagy ZP, Abdelmassih R, Balmaceda J. The effect of ultrasound (US) guided embryo transfer (ET) and the choice of catheter on the outcome of IVF. *Fertil Steril* 2001;**76**:S88–S89.
- Abu-Musa A, Usta I, Nassar A, Hajami F, Hannoun A. Effect of 17 $\alpha$ -hydroxyprogesterone caproate before embryo transfer on the outcome of in vitro fertilization and embryo transfer: a randomized trial. *Fertil Steril* 2008;**89**:1098–1102.
- Aflatoonian A, Ghandi S, Tabibnejad N. The effect of intercourse around embryo transfer on pregnancy rate in assisted reproductive technology cycles. *Int J Fertil Steril* 2009;**2**:169–72.
- Ahn J, Kim C, Kim S, Jeon G, Kim S, Chae H. Effects of administration of oxytocin antagonist on implantation and pregnancy rates in patients with repeated failure of IVF/ICSI treatment—clinical and experimental reproductive medicine. *Korean J Reprod Med* 2009;**36**:275–281.
- Allahbadia GN, Kadam K, Gandhi G, Arora S, Valliappan JB, Joshi A, Allahbadia S, Wolman I. Embryo transfer using the SureView catheter-beacon in the womb. *Fertil Steril* 2010;**93**:344–350.
- Al-Shawaf T, Dave R, Harper J, Linehan D, Riley P, Craft I. Transfer of embryos into the uterus: how much do technical factors affect pregnancy rates? *J Assist Reprod Genet* 1993;**10**:31–36.
- Amarin ZO, Obeidat BR. Bed rest versus free mobilisation following embryo transfer: a prospective randomised study. *BJOG* 2004;**111**:1273–1276.
- Ammar A, Mousa K, Hamed Rabei Galal NN. A. Effect of ultrasound guided embryo transfer on pregnancy rates. *New York Sci J* 2013;**6**:79–83.
- Amorcho B, Gomez E, Pontes L, Campos I, Landeras J, Munoz M, Ballesteros A, Remohí J, Pellicer A. Does the selection of catheter for embryo transfer affect the success rate of an ART unit? *Hum Reprod* 1999;**14**:205.
- Amui J, Check JH, Brasile D. Speculum retention during embryo transfer does not improve pregnancy rates following embryo transfer—a randomized study. *Clin Exp Obstet Gynecol* 2011;**38**:333–334.
- Andersen D, Løssl K, Andersen AN, Fürbringer J, Bach H, Simonsen J, Larsen EC. Acupuncture on the day of embryo transfer: a randomized controlled trial of 635 patients. *Reprod Biomed Online* 2010;**21**:366–372.

- Arefi S, Fazeli E, Esfahani M, Borhani N, Yamini N, Hosseini A, Farifteh F. Granulocyte-colony stimulating factor may improve pregnancy outcome in patients with history of unexplained recurrent implantation failure: an RCT. *Int J Reprod BioMed* 2018;**16**: 299–304.
- Arvas A, Karacan M, Ulug M, Cebi Z, Berberoglugil M, Batukan M, Camlibel T. A 20-second wait or immediate withdrawal of the catheter following ultrasound-guided blastocyst transfer. *Fertil Steril* 2014;**102**:e30.
- Azmy O, Taha T, Bibars M, Refaat M. Ultrasound guided versus blind touch embryo transfer: the Egyptian experience. In: *Proceedings of the 15th World Conference of IVF and 4th Congress of IVM* 2009, p.9–12. <http://www.edlearning.it/proceedings/moreinfo/20090419.htm> (October 2021, date last accessed).
- Balaban B, Urman B, Yakin K, Isiklar A, Kilic Y, Aksoy S. High pregnancy and implantation rates can be achieved in blastocyst transfers using hyaluronan enriched culture and transfer medium. *Fertil Steril* 2004;**82**:S221.
- Bar-Hava I, Meltzer S, Rabinson J, Ayash I, Segal S, Tur-Kaspa I. Ultrasound guided versus blind tactile embryo transfer: a prospective randomised study. *Hum Reprod* 2003;**18**:xviii7.
- Baris D, Ata AB, Isiklar A, Balaban B, Urman B. Prospective randomized comparison of Wallace and Labotect embryo transfer catheters. *Reprod Biomed Online* 2007;**14**:471–476.
- Benson MR, Elkind-Hirsch KE, Theall A, Fong K, Hogan RB, Scott RT. P-18: impact of acupuncture before and after embryo transfer on the outcome of in vitro fertilization cycles: a prospective single blind randomized study. *Fertil Steril* 2006;**86**:S135.
- Berkkanoglu M, Isikoglu M, Seleker M, Ozgur K. Flushing the endometrium prior to the embryo transfer does not affect the pregnancy rate. *Reprod Biomed Online* 2006;**13**:268–271.
- Bernabeu R, Roca M, Torres A, Ten J. Indomethacin effect on implantation rates in oocyte recipients. *Hum Reprod* 2006;**21**: 364–369.
- Bjuresten K, Hreinsson JG, Fridström M, Rosenlund B, Ek I, Hovatta O. Embryo transfer by midwife or gynecologist: a prospective randomized study. *Acta Obstet Gynecol Scand* 2003;**82**:462–466.
- Bodri D, Colodrón M, García D, Obradors A, Vernaev V, Coll O. Transvaginal versus transabdominal ultrasound guidance for embryo transfer in donor oocyte recipients: a randomized clinical trial. *Fertil Steril* 2011;**95**:2263–2268.e1.
- Boone WR, Johnson JE, Blackhurst DM, Crane MM. Short communication: Cook versus EdwardsWallace: are there differences in flexible catheters? *J Assist Reprod Genet* 2001;**18**:15–17.
- Bosch E, Aasted H, Klein BM, Arce J-C. A randomized, double-blind, placebo-controlled, multi-center, phase 2 trial to investigate the effect of barusiban on implantation in IVF/ICSI patients. *Fertil Steril* 2019;**112**:e179.
- Botta G, Grudzinskas G. Is a prolonged bed rest following embryo transfer useful? *Hum Reprod* 1997;**12**:2489–2492.
- Brook N, Khalaf Y, Coomarasamy A, Edgeworth J, Braude P. A randomized controlled trial of prophylactic antibiotics (co-amoxiclav) prior to embryo transfer. *Hum Reprod* 2006;**21**:2911–2915.
- Caanen MR, van der Houwen LE, Schats R, Vergouw CG, de Leeuw B, Lambers MJ, Groeneveld E, Lambalk CB, Hompes PG. Embryo transfer with controlled injection speed to increase pregnancy rates: a randomized controlled trial. *Gynecol Obstet Invest* 2016;**81**: 394–404.
- Cambiaghi AS, Leao RBF, Alvarez AV, Nascimento PF. Intrauterine injection of human chorionic gonadotropin before embryo transfer may improve clinical pregnancy and implantation rates in blastocysts transfers. *Fertil Steril* 2013;**100**:S121.
- Cardona Barberán A, Boel A, Vanden MF, Stoop D, Heindryckx B. Diagnosis and treatment of male infertility-related fertilization failure. *J Clin Med* 2020;**9**:3899.
- Catoire P, Delaunay L, Dannappel T, Baracchini D, Marcadet-Fredet S, Moreau O, Pacaud L, Przyrowski D, Marret E. Hypnosis versus diazepam for embryo transfer: a randomized controlled study. *Am J Clin Hypn* 2013;**55**:378–386.
- Chen S-C, Lai T-H, Lee F-K. The influence of abdominal ultrasound-guided embryo transfer on pregnancy rate: a preliminary report. *Fertil Steril* 2007;**87**:1235–1237.
- Cirillo F, Patrizio P, Baccini M, Morengi E, Ronchetti C, Cafaro L, Zannoni E, Baggiani A, Levi-Setti PE. The human factor: does the operator performing the embryo transfer significantly impact the cycle outcome? *Hum Reprod* 2020;**35**:275–282.
- Cornell J, Mulrow C, Localio R, Stack C, Meibohm A, Guallar E, Goodman S. Random-effects meta-analysis of inconsistent effects: a time for change. *Ann Intern Med* 2014;**160**:267–270.
- Coroleu B, Barri PN, Carreras O, Belil I, Buxaderas R, Veiga A, Balasch J. Effect of using an echogenic catheter for ultrasound-guided embryo transfer in an IVF programme: a prospective, randomized, controlled study. *Hum Reprod* 2006;**21**:1809–1815.
- Coroleu B, Barri PN, Carreras O, Martínez F, Parriego M, Hereter L, Parera N, Veiga A, Balasch J. The influence of the depth of embryo replacement into the uterine cavity on implantation rates after IVF: a controlled, ultrasound-guided study. *Hum Reprod* 2002a;**17**: 341–346.
- Coroleu B, Barri PN, Carreras O, Martínez F, Veiga A, Balasch J. The usefulness of ultrasound guidance in frozen-thawed embryo transfer: a prospective randomized clinical trial. *Hum Reprod* 2002b;**17**: 2885–2890.
- Coroleu B, Carreras O, Veiga A, Martell A, Martínez F, Belil I, Hereter L, Barri PN. Embryo transfer under ultrasound guidance improves pregnancy rates after in-vitro fertilization. *Hum Reprod* 2000;**15**:616–620.
- Cozzolino M, Vitagliano A, di Giovanni MV, Laganà AS, Vitale SG, Blaganje M, Drusany Starič K, Borut K, Patrelli TS, Noventa M. Ultrasound-guided embryo transfer: summary of the evidence and new perspectives. A systematic review and meta-analysis. *Reprod Biomed Online* 2018;**36**:524–542.
- Craig L, Rubin L, Peck J, Anderson M, Marshall L, Soules M. Acupuncture performed before and after embryo transfer: a randomized controlled trial. *J Reprod Med* 2014;**59**:313–320.
- Craig LB, Criniti AR, Hansen KR, Marshall LA, Soules MR. Acupuncture lowers pregnancy rates when performed before and after embryo transfer. *Fertil Steril* 2007;**88**:S40.
- Curfs MH, Cleine JH, van Kamp AA, Kruse-Blankestijn ME, Hondelink MN, Leerentveld RA. Comparison of the Wallace versus TDT embryo-transfer catheter: a prospective, randomized study. In *Third Biennial Alpha Conference* 2001. pp. 8–11.
- Dalal R, Pai H, Palshetkar N. Comparison of transabdominal (TA) versus transvaginally (TV) guided embryo transfer in oocyte recipient cycles: a prospective randomised trial. *BJOG* 2014;**121**:82.

- Davar R, Ghandi S, Tayebi N. Does transabdominal ultrasound-guided embryo transfer improve pregnancy rates in ART cycles? *Iran J Reprod Med* 2007;**5**:95–98.
- de Camargo Martins AMV, Baruffi RLR, Mauri AL, Petersen C, Oliveira JBA, Contart P, Pontes A, Franco JG. Ultrasound guidance is not necessary during easy embryo transfers. *J Assist Reprod Genet* 2004;**21**:421–425.
- Deep JP, Li Y, Liu H. The effect of embryo transfer on IVF treatment: comparing the trans-vaginal versus transabdominal ultrasound approach. *J Minimally Invasive Gynecol* 2013;**20**:S42–S43.
- Dehghani AS, Homayouni K, Kanannejad Z, Kanannejad Z. The effect of acupuncture on the day of embryo transfer on the in vitro fertilization outcomes: an RCT. *Int J Reprod Biomed* 2020;**18**:209–214.
- Dehghani Firouzabadi R, Ghandi S, Tayebi N. Effect of administration of single dose piroxicam before embryo transfer on implantation and pregnancy rates in IVF cycles. *J Biol Sci* 2006;**7**:123–126.
- Dehghani Firouzabadi R, Janati S, Razi MH. The effect of intrauterine human chorionic gonadotropin injection before embryo transfer on the implantation and pregnancy rate in infertile patients: a randomized clinical trial. *Int J Reprod Biomed* 2016;**14**:657–664.
- Devranoglu B, Özdamar Ö, Çakiroğlu Y, Küçükbaş M, Eken MK, Doğer E. The timing of embryo transfer catheter removal: should it be delayed or done immediately? A prospective randomized trial. *Gynecol Obstet Invest* 2018;**83**:29–34.
- Dittmann-Müller X, Zollner K, Zollner U. Prospective randomised clinical trial about the efficacy of a human embryo transfer medium (EmbryoGlue(R)). *Hum Reprod* 2009;**24**:167.
- Domar AD, Meshay I, Kelliher J, Alper M, Powers RD. The impact of acupuncture on in vitro fertilization outcome. *Fertil Steril* 2009;**91**:723–726.
- Drakeley AJ, Jorgensen A, Sklavounos J, Aust T, Gazvani R, Williamson P, Kingsland CR. A randomized controlled clinical trial of 2295 ultrasound-guided embryo transfers. *Hum Reprod* 2008;**23**:1101–1106.
- Duvan CI, Ozmen B, Satioglu H, Atabekoglu CS, Berker B. Does addition of low-dose aspirin and/or steroid as a standard treatment in nonselected intracytoplasmic sperm injection cycles improve in vitro fertilization success? A randomized, prospective, placebo-controlled study. *J Assist Reprod Genet* 2006;**23**:15–21.
- El-Shawarby SA, Ravhon A, Skull J, Ellenbogen A, Trew G, Lavery SA. Prospective randomized controlled trial of Wallace and Rocket embryo transfer catheters. *Reprod Biomed Online* 2008;**17**:549–552.
- El-Toukhy T, Sunkara SK, Khairy M, Dyer R, Khalaf Y, Coomarasamy A. A systematic review and meta-analysis of acupuncture in in vitro fertilisation. *BJOG* 2008;**115**:1203–1213.
- Eskandar M, Abou-Setta AM, Almushait MA, El-Amin M, Mohmad SEY. Ultrasound guidance during embryo transfer: a prospective, single-operator, randomized, controlled trial. *Fertil Steril* 2008;**90**:1187–1190.
- Eskandar M, Al-Emain M, Atwan Y, Bakar S. Does intrauterine injection of human chorionic gonadotropin before embryo transfer improve the pregnancy rate in vitro fertilization/intracytoplasmic sperm injection (IVF/ICSI) cycles? A prospective randomized controlled trial. *Reprod Sci* 2016;**23**:102A.
- Fancsovs P, Lehner A, Murber A, Kaszas Z, Rigo J, Urbancsek J. Effect of hyaluronan-enriched embryo transfer medium on IVF outcome: a prospective randomized clinical trial. *Arch Gynecol Obstet* 2015;**291**:1173–1179.
- Fancsovs P, Murber A, Gilan Z, Rigo J, Urbancsek J. Effect of hyaluronan containing transfer media on pregnancy and implantation rates in human IVF-ET cycles. A prospective randomized study. *Hum Reprod* 2011;**26**:124.
- Farquhar C. Introduction: add-ons for assisted reproductive technology: can we be honest here? *Fertil Steril* 2019;**112**:971–972.
- Fasano G, Antonacci R, Biramane J, Mbongolo G, Thi Vanhelleputte NM. C. Clinical outcomes after use of embryo-glue as a human embryo transfer (ET) medium in warming cycles. *Hum Reprod* 2016;**31**:204–205.
- Feichtinger W, Strohmer H, Radner KM, Goldin M. The use of fibrin sealant for embryo transfer: development and clinical studies. *Hum Reprod* 1992;**7**:890–893.
- Fekih M, Chachia S, Zarrouk W, Khairi H. Effect of ibuprophen administration before embryo transfer on IVF outcome: a randomized controlled trial. *Fertil Steril* 2013;**100**:S122.
- Foutouh I, Youssef M, Tolba M, Rushdi M, Nakieb A, Meguid W. Does embryo transfer catheter type affect pregnancy rate? *Middle East Fertil Soc* 2003;**8**:154–158.
- Franco JG, Martins AMVC, Baruffi RLR, Mauri AL, Petersen CG, Felipe V, Contart P, Pontes A, Oliveira JBA. Best site for embryo transfer: the upper or lower half of endometrial cavity? *Hum Reprod* 2004;**19**:1785–1790.
- Fratteirelli JL, Leondires MR, Fong K, Theall A, Locatelli S, Scott RT. Laser acupuncture before and after embryo transfer improves ART delivery rates: results of a prospective randomized double-blinded placebo controlled five-armed trial involving 1000 patients. *Fertil Steril* 2008;**90**:S105.
- Friedler S, Raziel A, Schachter M, Strassburger D, Kasterstein E, Komaroversusky D. Efficacy of hyaluronan-enriched embryo transfer medium in patients with repeated IVF-ET failures. *Hum Reprod* 2005;**20**:159.
- Friedler S, Schachter M, Strassburger D, Esther K, Ron El R, Raziel A. A randomized clinical trial comparing recombinant hyaluronan/recombinant albumin versus human tubal fluid for cleavage stage embryo transfer in patients with multiple IVF-embryo transfer failure. *Hum Reprod* 2007;**22**:2444–2448.
- Gaikwad S, Garrido N, Cobo A, Pellicer A, Remohi J. Bed rest after embryo transfer negatively affects in vitro fertilization: a randomized controlled clinical trial. *Fertil Steril* 2013;**100**:729–735.e2.
- Galiano V, Orvieto R, Machtinger R, Nahum R, Garzia E, Sulpizio P, Marconi AM, Seidman D. “Add-Ons” for assisted reproductive technology: do patients get honest information from fertility clinics’ websites? *Reprod Sci* 2021;**28**:3466–3472.
- García-Velasco JA, Isaza V, Martínez-Salazar J, Landazábal A, Requena A, Remohi J, Simón C. Transabdominal ultrasound-guided embryo transfer does not increase pregnancy rates in oocyte recipients. *Fertil Steril* 2002;**78**:534–539.
- García-Velasco JA, Martínez-Salazar J, Isaza V, Landazábal A, Requena A, Simón C. Does ultrasound guidance at embryo transfers improve pregnancy rates in oocyte recipients? *Fertil Steril* 2001;**76**:S75–S76.

- Gavrizi SZ, Skillern A. Does massage therapy immediately prior to embryo transfer improve clinical pregnancy rate in IVF-PGT-A (in-vitro fertilization-preimplantation genetic testing for aneuploidy) cycles? *Fertil Steril* 2019;**112**:e184–e185.
- Ghazzawi IM, Al-Hasani S, Karaki R, Sousa S. Transfer technique and catheter choice influence the incidence of transcervical embryo expulsion and the outcome of IVF. *Hum Reprod* 1999;**14**:677–682.
- Gilberto Almodin C, Cibebe V, Câmara M, Gonçalves P, Pereira C, Lopes Paixão C, Nakano R. Estudo comparativo entre dois cateteres diferentes usados para transferência de embriões [Comparative study between different catheters for embryo transfer]. n.d. JBRA Assist. Reprod 2009;13.4. Available online at: <http://materbaby.com.br/wp-content/uploads/2020/01/Artigo-Estudo-comparativo-entre-dois-cateteres-diferentes-usados-para-transfer%C3%Aancia-de-embri%C3%B5es.pdf> (October 2021, date last accessed).
- Glass KB, Green CA, Fluker MR, Schoolcraft WB, McNamee PI, Meldrum DR. Multicenter randomized controlled trial of cervical irrigation at the time of embryo transfer. *Fertil Steril* 2000;**74**:S31.
- GRADE Working Group. Grading quality of evidence and strength of recommendations. *BMJ* 2004;**328**:1490.
- Groutz A, Lessing JB, Wolf Y, Azem F, Yovel I, Amit A. Comparison of transmyometrial and transcervical embryo transfer in patients with previously failed in vitro fertilization-embryo transfer cycles and/or cervical stenosis. *Fertil Steril* 1997;**67**:1073–1076.
- Grunert GM, Dunn RC, Valdes CT, Wun CC, Wun WSA. Comparison of Wallace, Frydman DT and Cook embryo transfer catheter for IVF: a prospective randomised study [abstract]. *Fertil Steril* 1998;**70**:O–322.
- Gu YE, Zhang X, Zhang Q, Dai MC, Wu Y, Zhou Y, Qu F. The effects of acupuncture on pregnancy outcomes of in vitro fertilization with embryo transfer: an interdisciplinary systematic review. *J Gynecol Obstet Hum Reprod* 2019;**48**:677–684.
- Hafezi M, Madani T, Arabipour A, Zolfaghari Z, Sadeghi M, Ramezanali F. The effect of intrauterine human chorionic gonadotropin flushing on live birth rate after vitrified-warmed embryo transfer in programmed cycles: a randomized clinical trial. *Arch Gynecol Obstet* 2018;**297**:1571–1576.
- Hannoun A, Zreik TG, Ghaziri G, Abu Musa A, Awwad J. Effect of powdered gloves, worn at the time of embryo transfer, on the pregnancy outcome of IVF cycles. *J Assist Reprod Genet* 2009;**26**:25–27.
- Harbord R, Higgins J. METAREG: Stata module to perform meta-analysis regression. *Statistical Software Components*. Boston College Department of Economics, 2009. <https://ideas.repec.org/c/boc/bocode/s446201.html> (October 2021, date last accessed).
- Harbord RM, Harris RJ, Sterne JAC. Updated tests for small-study effects in meta-analyses. *Stata J* 2009;**9**:197–210.
- Hauzman E, Kohls G, Barrio A, Martinez-Salazar J, Iglesias C, Garcia-Velasco JA. Comparison of embryo transfer in egg donation recipients with transvaginal and transabdominal ultrasound: a randomized pilot study. *Hum Reprod* 2013;**28**:241.
- He Y, Wu H, He X, Xing Q, Zhou P, Cao Y, Wei Z. Administration of atosiban in patients with endometriosis undergoing frozen-thawed embryo transfer: a prospective, randomized study. *Fertil Steril* 2016;**106**:416–422.
- Hebisha SA, Aboelazm BA, Adel HM, Ahmed AI. Impact of the oxytocin receptor antagonist (ATOSIBAN) administered shortly before embryo transfer on pregnancy outcome after intracytoplasmic sperm injection (ICSI). *Fertil Steril* 2016;**106**:e88–e89.
- Hong KH, Forman EJ, Werner MD, Upham KM, Gumeny CL, Winslow AD, Kim TJ, Scott RT. Endometrial infusion of human chorionic gonadotropin at the time of blastocyst embryo transfer does not impact clinical outcomes: a randomized, double-blind, placebo-controlled trial. *Fertil Steril* 2014;**102**:1591–1595.e2.
- Hosseini RS, Farzad L, Abdollahi S, Nouri M, Ghasemzadeh A, Hamdi K, Soleimanpour H. Effect of intrauterine injection of human chorionic gonadotropin before frozen-thawed embryo transfer on implantation and clinical pregnancy rate: a randomized controlled trial. *Int J Womens Health Reprod Sci* 2016;**4**:200–203.
- Kan AKS, Abdalla HI, Gafar AH, Nappi L, Ogunyemi BO, Thomas A, Ola-Ojo OO. Embryo transfer: ultrasound-guided versus clinical touch. *Hum Reprod* 1999;**14**:1259–1261.
- Kandari S. Time lapse selected elective single embryo transfer in hyaluronan enriched transfer medium in PCOS improves live birth rates compared to use of conventional embryo transfer media. a possible alternative to freeze-all cycles in PCOS. *Fertil Steril* 2019;**112**:e47–e48.
- Karande V, Hazlett D, Vietzke M, Gleicher N. A prospective randomized comparison of the Wallace catheter and the Cook Echo-Tip® catheter for ultrasound-guided embryo transfer. *Fertil Steril* 2002;**77**:826–830.
- Karavani G, Ben-Meir A, Shufaro Y, Hyman JH, Revel A. Transvaginal ultrasound to guide embryo transfer: a randomized controlled trial. *Fertil Steril* 2017;**107**:1159–1165.
- Karimian L, Naghibi ZH, Yazdi PE, Moini A, Valojerd MR, Akhondi MM, Shahverdi AH, Movaghar B, Hassani F, Sedighi-Gilani MA. 147 The effect of intercourse on b HCG level and pregnancy outcome after IVF cycles retrospective evaluation of mechanical and laser assisted hatching efficiency in two different age groups. *Reprod Biomed Online* 2010;**20**:S65.
- Korosec S, Virant-Klun I, Tomazevic T, Zech NH, Meden-Vrtovec H. Single fresh and frozen-thawed blastocyst transfer using hyaluronan-rich transfer medium. *Reprod Biomed Online* 2007;**15**:701–707.
- Kosmas IP, Janssens R, Munck L, de Turki H, Al Elst J, van der Tournaye H, Devroey P. Ultrasound-guided embryo transfer does not offer any benefit in clinical outcome: a randomized controlled trial. *Hum Reprod* 2007;**22**:1327–1334.
- Krampl E, Zegermacher G, Eichler C, Obruca A, Strohmer H, Feichtinger W. Air in the uterine cavity after embryo transfer. *Fertil Steril* 1995;**63**:366–370.
- Laokirkkiat P, Thanaboonyawat I, Boonsuk S, Petyim S, Prechapanich J, Choavaratana R. Increased implantation rate after intrauterine infusion of a small volume of human chorionic gonadotropin at the time of embryo transfer: a randomized, double-blind controlled study. *Arch Gynecol Obstet* 2019;**299**:267–275.
- Lavery S, Ravhon A, Skull J, Ellenbogen A, Winston R. A prospective randomized controlled trial of Wallace and Rocket embryo transfer catheters in an IVF-embryo transfer programme [abstract]. *Hum Reprod* 2001;**16**:124.
- Leao R, Cambiaghi A, Leao B, Alvarez A, Figueiredo P. Intrauterine injection of human chorionic gonadotropin before embryo transfer



- may improve the pregnancy rates in in vitro fertilization cycles of patients with repeated implantation failures. In: *Proceedings of the 5th IVI International Congress*. 2013. Seville, Spain. Available online at: [https://www.comtecmed.com/ivi/2013/Uploads/Editor/abstract\\_66.pdf](https://www.comtecmed.com/ivi/2013/Uploads/Editor/abstract_66.pdf) (October 2021, date last accessed).
- Lensen S, Chen S, Goodman L, Rombauts L, Farquhar C, Hammarberg K. IVF add-ons in Australia and New Zealand: a systematic assessment of IVF clinic websites. *Aust N Z J Obstet Gynaecol* 2021;**61**:430–438.
- Levi-Setti PE, Albani E, Baggiani AM, Zannoni E, Colombo G, Liprandi V. Prospective randomized study comparing two soft catheters for embryo transfer. *Fertil Steril* 2002;**78**:S234–S235.
- Lewin A, Schenker JG, Avrech O, Shapira S, Safran A, Friedler S. The role of uterine straightening by passive bladder distension before embryo transfer in IVF cycles. *J Assist Reprod Genet* 1997;**14**:32–34.
- Li R, Lu L, Hao G, Zhong K, Cai Z, Wang W. Abdominal ultrasound-guided embryo transfer improves clinical pregnancy rates after in vitro fertilization: experiences from 330 clinical investigations. *J Assist Reprod Genet* 2005;**22**:3–8.
- Lin B, Pang Z, Jiang J. Fixed and random effects selection by REML and pathwise coordinate optimization. *J Comput Graph Stat* 2013;**22**:341–355.
- Lorusso F, Depalo R, Bettocchi S, Vacca M, Vimercati A, Selvaggi L. Outcome of in vitro fertilization after transabdominal ultrasound-assisted embryo transfer with a full or empty bladder. *Fertil Steril* 2005;**84**:1046–1048.
- MacLennan AH, Kerin JFP, Kirby C, Grant P, Warnes GM, Cox LW, Bryant-Greenwood G, Greenwood F. The effect of porcine relaxin vaginally applied at human embryo transfer in an in vitro fertilization programme. *Aust N Z J Obstet Gynaecol* 1985;**25**:68–71.
- Madani T, Ashrafi M, Jahangiri N, Abadi AB, Lankarani N. Improvement of pregnancy rate by modification of embryo transfer technique: a randomized clinical trial. *Fertil Steril* 2010;**94**:2424–2426.
- Madaschi C, Braga DPAF, de Cássia Savio Figueira R, Iaconelli A, Borges E. Effect of acupuncture on assisted reproduction treatment outcomes. *Acupunct Med* 2010;**28**:180–184.
- Mahani IM, Davar R. Hyaluronic acid versus albumin in human embryo transfer medium. *East Mediterr Health J* 2007;**13**:876–880.
- Mahlstedt PP, Macduff S, Bernstein J. Emotional factors and the in vitro fertilization and embryo transfer process. *J in Vitro Fert Embryo Transf* 1987;**4**:232–236.
- Mains L, van Voorhis BJ. Optimizing the technique of embryo transfer. *Fertil Steril* 2010;**94**:785–790.
- Maldonado LG, Ajzen SA, Busato WC, Iaconelli A, Bibancos M, Borges E. Impact of previous hysterosonometry on embryo transfer. *Fertil Steril* 2005;**84**:S364.
- Malhotra N, Sarkar P. Pregnancy outcome after bed rest versus early ambulation following embryo transfer during IVF/ICSI cycles—a randomised controlled study. *Fertil Steril* 2019;**112**:e146.
- Malina A, Pooley JA. Psychological consequences of IVF fertilization—review of research. *Ann Agric Environ Med* 2017;**24**:554–558.
- Mansour R. Minimizing embryo expulsion after embryo transfer: a randomized controlled study. *Hum Reprod* 2005;**20**:170–174.
- Mansour R, Tawab N, Kamal O, El-Faissal Y, Serour A, Aboulghar M, Serour G. Intrauterine injection of human chorionic gonadotropin before embryo transfer significantly improves the implantation and pregnancy rates in in vitro fertilization/intracytoplasmic sperm injection: a prospective randomized study. *Fertil Steril* 2011;**96**:1370–1374.e1.
- Marconi G, Young E, Vilela M, Bello A, Young E, Sueldo C. Prospective randomized comparison of an ultrasound-guided embryo transfer versus a blind catheter placement. *Fertil Steril* 2003;**80**:130.
- Martínez F, Coroleu B, Parriego M, Carreras O, Belil I, Parera N, Hereter L, Buxaderas R, Barri PN. Ultrasound-guided embryo transfer: immediate withdrawal of the catheter versus a 30 second wait. *Hum Reprod* 2001;**16**:871–874.
- Matorras R, Urquijo E, Mendoza R, Corcóstequi B, Expósito A, Rodríguez-Escudero F. Ultrasound-guided embryo transfer improves pregnancy rates and increases the frequency of easy transfers. *Hum Reprod* 2002;**17**:1762–1766.
- Mayer JF, Nehchiri F, Jones EC, Weedon VM, Kalin HL, Lanzendorf SE, Oehinger SC, Toner JP, Muasher SJ. Prospective randomized analysis of the impact of two different transfer catheters on clinical pregnancy rates [abstract]. *Fertil Steril* 1999;**1**:144–145.
- McDonald JA, Norman RA. Randomized controlled trial of a soft double lumen embryo transfer catheter versus a firm single lumen catheter: significant improvements in pregnancy rates. *Hum Reprod* 2002;**17**:1502–1506.
- McIlveen M, Lok FD, Pritchard J, Lashen H. Modern embryo transfer catheters and pregnancy outcome: a prospective randomized trial. *Fertil Steril* 2005;**84**:996–1000.
- Meriano J, Weissman A, Greenblatt EM, Ward S, Casper RF. The choice of embryo transfer catheter affects embryo implantation after IVF. *Fertil Steril* 2000;**74**:678–682.
- Mitchell JD, Wardle PG, Foster PA, Hull MGR. Effect of bladder filling on embryo transfer. *J In Vitro Fert Embryo Transf* 1989;**6**:263–265.
- Moini A, Kiani K, Bahmanabadi A, Akhoond M, Akhlaghi A. Improvement in pregnancy rate by removal of cervical discharge prior to embryo transfer in ICSI cycles: a randomised clinical trial. *Aust N Z J Obstet Gynaecol* 2011;**51**:315–320.
- Moon HS, Park SH, Lee JO, Kim KS, Joo BS. Treatment with piroxicam before embryo transfer increases the pregnancy rate after in vitro fertilization and embryo transfer. *Fertil Steril* 2004;**82**:816–820.
- Moraga-Sanchez R, Saucedo-De La Llata E, Batiza-Resendiz V, Santos-Haliscak R, Galache-Vega P. Ultrasound influence and 30 second wait in embryo transfer. *Hum Reprod* 2004;**19**:127.
- Moraloglu O, Tonguc E, Var T, Zeyrek T, Batioglu S. Treatment with oxytocin antagonists before embryo transfer may increase implantation rates after IVF. *Reprod Biomed Online* 2010;**21**:338–343.
- Morbeck D. Prospective randomized clinical trial of novel implantation promoting medium (EmbryoGlue) to improve IVF success rates [NCT00588250] ClinicalTrials.gov. 2007. Available at: <https://www.clinicaltrials.gov/ct2/show/NCT00588250> (Date last accessed October 2021).
- Moreno V, Balasch J, Vidal E, Calafell JM, Cívico S, Vanrell JA. Air in the transfer catheter does not affect the success of embryo transfer. *Fertil Steril* 2004;**81**:1366–1370.
- Mortimer ST, Fluker M, Yuzpe A. Effect of embryo transfer catheter on implantation rates. *Fertil Steril* 2002;**78**:S17–S18.



- Mostajeran F, Godazandeh F, Ahmadi SM, Movahedi M, Jabalamelian SA. Effect of intrauterine injection of human chorionic gonadotropin before embryo transfer on pregnancy rate: a prospective randomized study. *J Res Med Sci* 2017;**22**:6.
- Moy I, Milad M, Barnes R, Confino E, Kazer R, Zhang X. Randomized controlled trial: effects of acupuncture on pregnancy rates in women undergoing in vitro fertilization. *Fertil Steril* 2011;**95**:583–587.
- Murphy EM, Nichols J, Somkuti SG, Sobel M, Braverman A, Barmat LI. Randomized trial of harp therapy during in vitro fertilization-embryo transfer. *J Evid Based Complementary Altern Med* 2014;**19**: 93–98.
- Nancarrow L, Tempest N, Drakeley AJ, Homburg R, Russell R, Hapangama DK. National survey highlights the urgent need for standardisation of embryo transfer techniques in the UK. *J Clin Med* 2021;**10**:2839.
- Nazari A, Askari HA, Check JH, O'Shaughnessy A. Embryo transfer technique as a cause of ectopic pregnancy in in vitro fertilization. *Fertil Steril* 1993;**60**:919–921.
- Ng EHY, Li RHW, Chen L, Lan VTN, Tuong HM, Quan S. A randomized double blind comparison of atosiban in patients undergoing IVF treatment. *Hum Reprod* 2014;**29**:2687–2694.
- Ng KKL, Rozen G, Stewart T, Agresta F, Polyakov A. Does nifedipine improve outcomes of embryo transfer? Interim analysis of a randomized, double blinded, placebo-controlled trial. *Medicine (Baltimore)* 2019;**98**:e14251.
- Nihat Candan Z, Avcil F, Özden H, Uslu H, Karaman Y. Comparing the performance of the Cook and the Prodimed embryo transfer catheters. *Reprod Biomed Online* 2014;**28**:S15.
- Obidniak D, Gzgyan A, Dzhelekhianova L, Feoktistov A. Effect of colony-stimulating growth factor on outcome of frozen-thawed embryo transfer in patients with repeated implantation failure. *Fertil Steril* 2016;**106**:e134–e135.
- Obidniak D, Gzgyan A, Feoktistov A, Niauri D. Randomized controlled trial evaluating efficacy of autologous platelet-rich plasma therapy for patients with recurrent implantation failure. *Fertil Steril* 2017;**108**:e370.
- Ocal P, Cepni I, Idil MH, Salihoglu F, Irez T, Aksu F. Appropriate embryo transfer technique improves the pregnancy rate in human in vitro fertilization. *Middle East Fertil Soc J* 2003;**8**:65–68.
- Omodei U, Piccioni G, Tombesi S, Dordoni D, Fallo L, Ghilardi F. Effect of acupuncture on rates of pregnancy among women undergoing in vitro fertilization. *Fertil Steril* 2010;**94**:266.
- Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, Shamseer L, Tetzlaff JM, Akl EA, Brennan SE. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ* 2021;**372**.
- Paulus WV, Zhang M, Strehler E, El-Danasouri I, Sterzik K. Influence of acupuncture on the pregnancy rate in patients who undergo assisted reproduction therapy. *Fertil Steril* 2002;**77**:721–724.
- Paulus WV, Zhang M, Strehler E, Seybold B, Sterzik K. Placebo-controlled trial of acupuncture effects in assisted reproduction therapy. *Hum Reprod* 2003;**18**:18–19.
- Peikrishvili R, Evrard B, Pouly JL, Janny L. L'antibiothérapie prophylactique (amoxicilline + acide clavulanique) avant transfert pour fécondation in vitro est inutile: résultats d'une étude randomisée. *J Gynécol Obstét Biol Reprod* 2004;**33**:713–719.
- Penzias A, Bendikson K, Butts S, Coutifaris C, Falcone T, Fossum G, Gitlin S, Gracia C, Hansen K, La Barbera A et al. Performing the embryo transfer: a guideline. *Fertil Steril* 2017a;**107**:882–896.
- Penzias A, Bendikson K, Butts S, Coutifaris C, Falcone T, Fossum G, Gitlin S, Gracia C, Hansen K, Mersereau J et al.; Practice Committee of the American Society for Reproductive Medicine. Electronic address: ASRM@asrm.org. Standard embryo transfer protocol template: a committee opinion. *Fertil Steril* 2017b;**107**: 897–900.
- Perez O, Adriaanse H, Tilley B, Navarrete G, Lay L, Little LM, Gada R, Lawrence L, Lee K, Thomas MR. The effect of extended blastocyst exposure of hyaluronan enriched transfer media on implantation rate in frozen embryo transfers. *Fertil Steril* 2019;**112**:e438.
- Perin PM, Neves PA, Maluf M. The influence of two different transfer catheters on the pregnancy rate in a human in vitro fertilization/embryo transfer program. *Fertil Steril* 1997;**1997**:220.
- Porat N, Boehnlein LM, Schouweiler CM, Kang J, Lindheim SR. Interim analysis of a randomized clinical trial comparing abdominal versus transvaginal ultrasound-guided embryo transfer. *J Obstet Gynaecol Res* 2010;**36**:384–392.
- Pottie K, Gorber SC, Singh H, Joffres M, Lindsay P, Brauer P, Jaramillo A, Tonelli M. Estimating benefits and harms of screening across subgroups: the Canadian Task Force on Preventive Health Care integrates the GRADE approach and overcomes minor challenges. *J Clin Epidemiol* 2012;**65**:1245–1248.
- Prapas Y, Prapas N, Hatziparasidou A, Prapa S, Nijs M, Vanderzwalmen P, Vlassis G, Jones EE. The echoguide embryo transfer maximizes the IVF results. *Acta Eur Fertil* 1995;**26**: 113–115.
- Prato LD, Borini A. Effect of piroxicam administration before embryo transfer on IVF outcome: a randomized controlled trial. *Reprod Biomed Online* 2009;**19**:604–609.
- Purcell KJ, Schembri M, Telles TL, Fujimoto VY, Cedars MI. Bed rest after embryo transfer: a randomized controlled trial. *Fertil Steril* 2007;**87**:1322–1326.
- Qu F, Zhang D, Chen L-T, Wang F-F, Pan J-X, Zhu Y-M, Ma C-M, Huang Y-T, Ye X-Q, Sun S-J et al. Auricular acupressure reduces anxiety levels and improves outcomes of in vitro fertilization: a prospective, randomized and controlled study. *Sci Rep* 2014;**4**: 5028.
- Ravhon A, Nahum H, Weissman A, Biran G, Umansky N, Levran D. Embryo transfer in hyaluronan enriched transfer medium does not improve pregnancy rate in IVF treatment. *Fertil Steril* 2005;**84**: S376–S377.
- Revelli A, Rovei V, Dalmasso P, Gennarelli G, Racca C, Evangelista F, Benedetto C. Large randomized trial comparing transabdominal ultrasound-guided embryo transfer with a technique based on uterine length measurement before embryo transfer. *Ultrasound Obstet Gynecol* 2016;**48**:289–295.
- Rezábek K, Koryntova D, Zivny J. Does bedrest after embryo transfer cause a worse outcome in in vitro fertilization? *Ceska Gynekol* 2001;**66**:175–178.
- Rhodes TL, Higdon HL, Boone WR. Comparison of pregnancy rates for two embryo-transfer catheters. *Fertil Steril* 2007;**87**:411–416.
- Rinaldi L, Floccari A, Selman H. Ultrasound guidance of embryo transfer: a role for midwife. *Sex Reprod Healthc* 2014;**5**:47–49.

- Ruhlman C, Bisoli C, Terrado G, Rolla ED, Nicholson RE, Gnocchi D. The inefficacy of cervical mucus aspiration prior to embryo transfer: a prospective randomized trial [abstract]. ASRM/CFAS Conjoint Annual Meeting. *Fertil Steril* 1999;**71**:154.
- Saldeen P, Abou-Setta AM, Bergh T, Sundström P, Holte J. A prospective randomized controlled trial comparing two embryo transfer catheters in an ART program. *Fertil Steril* 2008;**90**:599–603.
- Sallam HN, Agameya AF, Rahman AF, Ezzeldin F, Sallam AN. Ultrasound measurement of the uterocervical angle before embryo transfer: a prospective controlled study. *Hum Reprod* 2002;**17**:1767–1772.
- Santibañez Á, García J, Pashkova O, Colín O, Castellanos G, Sánchez AP, de la Jara JF. Effect of intrauterine injection of human chorionic gonadotropin before embryo transfer on clinical pregnancy rates from in vitro fertilisation cycles: a prospective study. *Reprod Biol Endocrinol* 2014;**12**:9.
- Saravelos SH, Kong GWS, Chung JPW, Mak JSM, Chung CHS, Cheung LP, Li T-C. A prospective randomized controlled trial of 3D versus 2D ultrasound-guided embryo transfer in women undergoing ART treatment. *Hum Reprod* 2016;**31**:2255–2260.
- Schoolcraft W, Lane M, Stevens J, Gardner DK. Increased hyaluronan concentration in the embryo transfer medium results in a significant increase in human embryo implantation rate. *Fertil Steril* 2002;**78**:S5.
- Schwarze JE, Crosby J, Mackenna A. Atosiban improves the outcome of embryo transfer. A systematic review and meta-analysis of randomized and non-randomized trials. *JBRA Assist Reprod* 2020;**24**:421–427.
- Seto MTY, Cheung KW, Lo TK, Ng EHY. Pregnancy outcomes of women randomized to receive real versus placebo acupuncture on the day of fresh or frozen-thawed embryo transfer. *Eur J Obstet Gynecol Reprod Biol* 2017;**218**:119–122.
- Shaker AG, Fleming R, Jamieson ME, Yates RWS, Coutts JRT. Infertility: assessments of embryo transfer after in-vitro fertilization: effects of glyceryl trinitrate. *Hum Reprod* 1993;**8**:1426–1428.
- Sigalos GA, Michalopoulos Y, Kastoras AG, Triantafyllidou O, Vlahos NF. Low versus high volume of culture medium during embryo transfer: a randomized clinical trial. *J Assist Reprod Genet* 2018;**35**:693–699.
- Simon A, Safran A, Revel A, Aizenman E, Reubinoff B, Porat-Katz A, Lewin A, Laufer N. Hyaluronic acid can successfully replace albumin as the sole macromolecule in a human embryo transfer medium. *Fertil Steril* 2003;**79**:1434–1438.
- Singh R, Singh M. Intra-uterine administration of human chorionic gonadotrophin (hCG) before embryo transfer in recurrent implantation failure (RIF) patients improves implantation and pregnancy rates in IVF-ICSI cycles. *Hum Reprod* 2014;**29**:79.
- Singh R, Singh M. P-296 RCT of intra-uterine administration or subcutaneous injection of GCSF (granulocyte colony-stimulating factor) before embryo-transfer on resistant thin endometrium in IVF. *Hum Reprod* 2018;**33**:i1–i541.
- Singh R, Singh M, Jindal A, Jindal P. A prospective randomized controlled study (RCT) of intra-uterine administration of granulocyte colony-stimulating factor (G-CSF) before embryo-transfer on resistant thin endometrium in IVF cycles. *Hum Reprod* 2015;**30**:120.
- Smith CA, Armour M, Shewamene Z, Tan HY, Norman RJ, Johnson NP. Acupuncture performed around the time of embryo transfer: a systematic review and meta-analysis. *Reprod Biomed Online* 2019;**38**:364–379.
- So E, Ng E, Wong Y, Lau E, Yeung W, Ho P. A randomized double blind comparison of real and placebo acupuncture in IVF treatment. *Hum Reprod* 2009;**24**:341–348.
- So E, Ng E, Wong Y, Yeung W, Ho P. Acupuncture for frozen-thawed embryo transfer cycles: a double-blind randomized controlled trial. *Reprod Biomed Online* 2010;**20**:814–821.
- Song Z-R, Zhao X-H, Bai X-H, Lu Y-H, Zhang H-J, Wang Y-X, Lu R. Application of oxytocin antagonists in thaw embryo transfer. *Zhonghua Fu Chan Ke Za Zhi* 2013;**48**:667–670.
- Soroka D, Wells G, Kotaba DD. Does the aspiration of cervical mucus prior to embryo transfer in women undergoing in vitro fertilization and embryo transfer improve pregnancy rates? *Fertil Steril* 1999;**72**:S102.
- Steptoe PC, Edwards RG. Birth after the reimplantation of a human embryo. *Lancet* 1978;**2**:366.
- Sterne JAC, Harbord RM. Funnel plots in meta-analysis. *Stata J* 2004;**4**:127–141.
- Sterne JAC, Savović J, Page MJ, Elbers RG, Blencowe NS, Boutron I, Cates CJ, Cheng HY, Corbett MS, Eldridge SM. RoB 2: a revised tool for assessing risk of bias in randomised trials. *BMJ* 2019;**366**:14898.
- Stocker LJ, Hardingham KL, Cheong YC. A randomized controlled trial assessing whether listening to music at time of embryo transfer effects anxiety levels. *Gynecol Obstet* 2016;**6**:1–6.
- Sutton AJ, Duval SJ, Tweedie RL, Abrams KR, Jones DR. Empirical assessment of effect of publication bias on meta-analyses. *BMJ* 2000;**320**:1574–1577.
- Talwar P, Naredi N, Sandeep K, Joneja G, Duggal B. Does catheter choice during embryo transfer alter the pregnancy rate? *Med J Armed Forces India* 2011;**67**:311–314.
- Tang OS, Ng EHY, So WWK, Ho PC. Ultrasound-guided embryo transfer: a prospective randomized controlled trial. *Hum Reprod* 2001;**16**:2310–2315.
- Teixeira DM, Dassuncão LA, Vieira CVR, Barbosa MAP, Coelho Neto MA, Nastri CO, Martins WP. Ultrasound guidance during embryo transfer: a systematic review and meta-analysis of randomized controlled trials. *Ultrasound Obstet Gynecol* 2015;**45**:139–148.
- Tierney JF, Pignon JP, Gueffier F, Clarke M, Askie L, Vale CL, Burdett S; Cochrane IPD Meta-analysis Methods Group. How individual participant data meta-analyses have influenced trial design, conduct, and analysis. *J Clin Epidemiol* 2015;**68**:1325–1335.
- Tomari H, Honjou K, Kunitake K, Nishimura K, Hidaka N, Nagata Y. Effect of embryo glue transfer medium during fresh and frozen-thawed embryo transfer. *Fertil Steril* 2014;**102**:e313.
- Urman B, Yakin K, Ata B, Isiklar A, Balaban B. Effect of hyaluronan-enriched transfer medium on implantation and pregnancy rates after day 3 and day 5 embryo transfers: a prospective randomized study. *Fertil Steril* 2008;**90**:604–612.
- Valojerdi MR, Karimian L, Yazdi PE, Gilani MAS, Madani T, Baghestani AR. Efficacy of a human embryo transfer medium: a prospective, randomized clinical trial study. *J Assist Reprod Genet* 2006;**23**:207–212.
- Visnova H, Tournaye HJ, Humberstone A, Terrill P, Macgregor L, Loumaye E. A placebo-controlled, randomized, double-blind, phase 3 study assessing ongoing pregnancy rates after single oral

- administration of a novel oxytocin receptor antagonist, nolasiban, prior to single embryo transfer. *Fertil Steril* 2018;**110**:e45.
- Visschers B, Bots R, Peeters M, Mol B, van Dessel H. Removal of cervical mucus: effect on pregnancy rates in IVF/ICSI. *Reprod Biomed Online* 2007;**15**:310–315.
- Walker DL, Thornhill AR, Allemand MC, Tatpati LL, Wentworth MA, Tummon IS. A randomized, controlled, double blinded trial of EmbryoGlue® in frozen embryo transfer cycles: an interim analysis. *Fertil Steril* 2005;**84**:S415–S416.
- Wang M, Deng H, Ye H. Intrauterine injection of human chorionic gonadotropin improves pregnancy outcome in patients with repeated implantation failure in frozen-thawed embryo transfer. *Zhong Nan Da Xue Xue Bao Yi Xue Ban* 2019;**44**:1247–1251.
- van Weering HGI, Schats R, McDonnell J, Vink JM, Vermeiden JPW, Hompes PGA. The impact of the embryo transfer catheter on the pregnancy rate in IVF. *Hum Reprod* 2002;**17**:666–670.
- Weissman A, Farhi J, Steinfeld Z, Mutsafi R, Nahum H, Levran D. A prospective, randomized study of ultrasound-guided embryo transfer. *Fertil Steril* 2003;**80**:122.
- Westergaard L, Mao Q, Krogslund M, Sandrini S, Lenz S, Grinsted J. Acupuncture on the day of embryo transfer significantly improves the reproductive outcome in infertile women: a prospective, randomized trial. *Fertil Steril* 2006;**85**:1341–1346.
- Wirleitner B, Schuff M, Vanderzwalmen P, Stecher A. O-182 The usefulness of intrauterine hCG administration prior to blastocyst transfer in IVF-patients  $\geq 38$  years. *Hum Reprod* 2015a;**30**:i1–i501.
- Wirleitner B, Schuff M, Vanderzwalmen P, Stecher A, Okhowat J, Hradecký L, Kohoutek T, Králíková M, Spitzer D, Zech NH. Intrauterine administration of human chorionic gonadotropin does not improve pregnancy and live birth rates independently of blastocyst quality: a randomised prospective study. *Reprod Biol Endocrinol* 2015b;**13**:70.
- Wisanto A, Janssens R, Deschacht J, Camus M, Devroey P, van Steirteghem AC. Performance of different embryo transfer catheters in a human in vitro fertilization program. *Fertil Steril* 1989;**52**:79–84.
- Yakin K, Balaban B, Isiklar A, Bozdog H, Urman B. Improved clinical outcome in frozen-thawed embryo transfers with the use of hyaluronan-enriched medium. *Hum Reprod* 2004;**19**:89.
- Yao Z, Vansteelandt S, Elst J, van der Coetsier T, Dhont M, de Sutter P. The efficacy of the embryo transfer catheter in IVF and ICSI is operator-dependent: a randomized clinical trial. *Hum Reprod* 2009a;**24**:880–887.
- Yao Z, Vansteelandt S, Elst J, van der Coetsier T, Dhont M, de Sutter P. The efficacy of the embryo transfer catheter in IVF and ICSI is operator-dependent: a randomized clinical trial. *Hum Reprod* 2009b;**24**:880–887.
- Yayla Abide C, Ozkaya E, Sanverdi I, Bostancı Ergen E, Kurek Eken M, Devranoglu B, Bilgiç BE, Kilicci C, Kayatas Eser S. Prospective randomized trial comparing embryo transfers of cases with and without catheter rotation during its withdrawal. *Gynecol Obstet Invest* 2018;**83**:397–403.
- Yuan C, Song H, Fan L, Su S, Dong B. The effect of atosiban on patients with difficult embryo transfers undergoing in vitro fertilization—embryo transfer. *Reprod Sci* 2019;**26**:1613–1617.
- Yung SSF, Lai SF, Lam MT, Lui EMW, Ko JKY, Li HWR, Wong JYY, Lau EYL, Yeung WSB, Ng EHY. Hyaluronic acid-enriched transfer medium for frozen embryo transfer: a randomized, double-blind, controlled trial. *Fertil Steril* 2021;**116**:1001–1009.
- Zarei A, Homayoon N, Hessami K, Hashemi A, Davoodi S, Razavi B, Ghasempour L, Alamdarloo SM. Effect of piroxicam administration on outcome of frozen-thawed embryo transfer: a randomized, double-blinded, placebo-controlled trial. *J Obstet Gynaecol Res* 2021;**47**:296–301.
- Zarei A, Parsanezhad ME, Younesi M, Alborzi S, Zolghadri J, Samsami A, Amooee S, Aramesh S. Intrauterine administration of recombinant human chorionic gonadotropin before embryo transfer on outcome of in vitro fertilization/intracytoplasmic sperm injection: a randomized clinical trial. *Iran J Reprod Med* 2014;**12**:1–6.
- Zegers-Hochschild F, World Health Organization, Adamson GD, Mouzon J, de Ishihara O, Mansour R, Nygren K, Sullivan E, Vanderpoel S. International Committee for Monitoring Assisted Reproductive Technology (ICMART) and the World Health Organization (WHO) revised glossary of ART terminology, 2009. *Fertil Steril* 2009;**92**:1520–1524.
- Zemyarska MS. Is it ethical to provide IVF add-ons when there is no evidence of a benefit if the patient requests it? *J Med Ethics* 2019;**45**:346–350.
- Zhang R, Feng X-J, Guan Q, Cui W, Zheng Y, Sun W, Han J-S. Increase of success rate for women undergoing embryo transfer by transcutaneous electrical acupoint stimulation: a prospective randomized placebo-controlled study. *Fertil Steril* 2011;**96**:912–916.
- Zhong J, Zhang L. [Transcutaneous electrical acupoint stimulation for pregnancy of in vitro fertilization-embryo transfer]. *Zhongguo Zhen Jiu* 2017;**37**:253–255.